

Interactive comment on “Mars MOURA magnetometer demonstration for high resolution mapping on terrestrial analogues” by M. Diaz Michelena et al.

M. Diaz Michelena et al.

diazma@inta.es

Received and published: 18 March 2016

Dear Prof. Martín,

We have read through your comments and have modified the manuscript accordingly.
(Text attached with changes highlighted: changes in yellow concern your comments)

We hope that the modifications accomplish with your expectation.

Sincerely, Marina

Mars MOURA magnetometer demonstration for high resolution mapping on terrestrial analogues

[Printer-friendly version](#)

[Discussion paper](#)



Marina Diaz-Michelena¹, Rolf Kilian^{2,3}, Ruy Sanz¹, Francisco Rios², Oscar Baeza²

¹ Payloads and Space Sciences Department, INTA, Ctra. Torrejón – Ajalvir km 4, 28850 Torrejón de Ardoz, Spain ² Geology Department, University of Trier, Behringstrasse, 54286 Trier, Germany ³ University of Magallanes, Punta Arenas, Chile

Abstract Satellite-based magnetic measurements of Mars indicate complex and very strong magnetic anomalies, which led to an intensive and long-lasting discussion about their possible origin. To make some progress in the investigation of the origin of these anomalies MOURA vector magnetometer was developed for in situ measurements on Mars. In this work we propose the utilisation of such instrument for future planetary on ground surveys. The proof of its suitability is done by testing on various terrestrial analogues characterised by most distinct magnetic anomalies of their basement rocks: (1) A magnetite body of EL Laco (up to + 110,000 nT) and its transition to surrounding andesites (< + 2,000 nT) in the Northern Andes of Chile showing the highest local magnetic anomalies. The magnetite-bearing ore body has highly variable local anomalies due to their complex formation history where a significant dispersion in paleo-orientations has been previously reported, while our vector data show relatively uniform and probably induced declinations. (2) A basaltic spatter cone of the Pali Aike volcanic field, in Southern Chile, was characterised by very strong magnetic anomalies along the crater rim (up to + 12,000 nT), controlled by the amount of single domain magnetites in the ground mass of the basalts. Due to their strong remanent signature paleodeclinations of the lavas and reorientations of collapsed blocks could be constrained by the vector data. (3) The Monturaqui meteorite crater (350 m diameter), in Northern Chile, shows significant variations of its anomalies (from - 2,000 to > + 6,000 nT) in restricted areas of several square metres along its crater rim related to unexposed iron-bearing fragments of the impactor while its granitic and ignimbritic target rocks exhibit only very weak anomalies. (4) An area with several amphibolitic dykes which cross-cut a Cretaceous granitoid in the southernmost Andes, where a decimetre-scale mapping was performed. In this case, pyrrhotite is the only magnetic

carrier. It was formed during hydrothermal processes within the dykes. Very low (+ 40 to + 120 nT) positive magnetic anomalies clearly depict the amount of 1 to 4 vol.% pyrrhotite in these dykes, which is important as a mineralogical indicator as well as to detect associated gold and copper enrichment.

(355 words)

Keywords: Andes, El Laco, Monturaqui, Pali Aike, Patagonian Batholith, continental crust, martian crust, granite, basalt, mafic dykes, volcanic crater, magnetic anomalies, magnetic carrier, magnetic susceptibility, ground magnetic survey, caesium magnetometer, MOURA magnetometer, Koenigsberger ratios, plutonic rocks, impact crater, iron meteorite, pyrrhotite, magnetite, hydrothermal mineralization, gold, copper. 1. Introduction

Mars magnetic field has been exhaustively measured between 100 and 440 km altitude by Mars Global Surveyor (Acuña et al., 1998; Connerney et al., 2005; Morschhauser et al., 2014). These data show that the magnetic anomalies of the martian crust are up to 20 times higher than those of the Earth (Scott and Fuller, 2004). However, a profound understanding of the magnetic signature of the martian crust would require mapping at different altitudes and consequently with a different magnetic zoom apart from further petrological analyses. Since Mars presents a very low dense atmosphere, aeromagnetic surveys are not achievable in the short term. Thus, on ground magnetometry with landers and rovers seems to be the most immediate feasible technology to complement the satellite measurements. MOURA magnetometer was developed by INTA in the context of MetNet Precursor Mission to perform vector magnetometry and gradiometry during on ground prior to rover-based surveys on extra-terrestrial planets, like Mars. The instrument has a very low mass (72 g), a scalable range, high precision, low detectable fields and noise, and is capable to work in the very hard environmental conditions of Mars. Diaz-Michelena et al. (2015 a) describes MOURA main technical details, the calibration, as well as the demonstration to measure the absolute magnetic field and its temporal variations. The first objective of the present work is

[Printer-friendly version](#)

[Discussion paper](#)



to demonstrate the capability of the miniaturized MOURA instrument in a real context of terrestrial analogues surveys by means of the inter-comparison with the data of a scalar caesium reference magnetometer (Diaz-Michelena and Kilian 2013). To do this, four different sites with a wide variability in the intensity of their magnetic signatures have been selected. A second objective is the magnetic investigation of these sites and their implication as terrestrial analogues of Mars. A further objective is the potential of high resolution mapping to show the lowest magnetic contrasts in the terrain and their correlation with the distinct magnetic carriers responsible for their signatures (Acuña et al., 1998; Connerney et al., 2005; Lillis et al., 2013). Our selected sites include magnetic anomalies from complex geological environments where e.g. noise factors, geometrical characteristics of non-exposed rock units and effects of the terrain relief can obscure partly the interpretations concerning the kind and magnetic effects of non-exposed rock units. Thus a modelling of the magnetic anomalies is desirable in general to improve the interpretation of geometrical and compositional effects of non-exposed rocks (e.g. Eppelbaum et al., 2015; Eppelbaum and Mishne, 2011; Jalongo et al., 2014). Since we are aware of this problematics, primarily we focus on the interpretation of the effects of distinct types and compositions of exposed rocks concerning the observed magnetic anomalies. Future more detailed studies of the investigated sites should include the above mentioned magnetic modelling which is out of the scope of this magnetometer demonstration.

2. Methodology

2.1. Magnetic Instrumentation

Two different magnetometers have been used for the present surveys: a conventional caesium scalar magnetometer: model G-858 MagMapper by Geometrics and MOURA vector magnetometer designed and developed by INTA MetNet team for Mars exploration. G-858 is taken as the reference magnetometer because it is a well-established hand-held instrument (8 - 9 kg) for magnetic surveys (ordnance, archaeology, environmental, mineralogy and petroleum prospection). It has 8 hours autonomy, provides

[Printer-friendly version](#)[Discussion paper](#)

a suitable contrast related to magnetic anomalies ($8 \text{ pT}/\sqrt{\text{Hz}}$) and good stability covering the range of the Earth magnetic field with a dynamic range between 20,000 to 100,000 nT. It also has several modes of operation: continuous and discrete to allow users to plan the prospections grids. In contrast to MOURA, this magnetometer only provides the intensity of the total magnetic field, its performance is dependent on the orientation of the head respect to the field, which changes significantly in the latitude range of the present survey, and it is restricted to areas with gradients higher than 20,000 nT/m (Table 1). MOURA is a vector magnetometer with two 3-axes magnetic sensors of Anisotropic MagnetoResistance (AMR) by Honeywell to build up a compact and miniaturized instrument (72 g mass and 67.5 cm³) for Mars exploration. The power consumption is limited to 400 - 430 mW, so it can operate during more than 10 hours with commercial batteries, and insignificant increase of weight to the user (3 x 25 g). The instrument is also designed for continuous or discrete modes of operation, it can work in every orientation with no incidence in the performance, and it is practically immune to gradients due to the small size of the transducer (μm -size). The characteristics are designed for Mars surface environment (- 90 to + 20 °C in operation, - 120 to + 125 °C in storage, and a total irradiance dose of 15 krad/s). The resolution is limited by the transducer (0.2 nT) and the range is adapted to that of the Earth geomagnetic field $\pm 65,000 \text{ nT}$ with an extended range in the auto-offset compensation mode of $\pm 130,000 \text{ nT}$ (Table 1, Diaz-Michelena et al., 2015 a).

2.2 Track performance

The tracks have been defined to cover most of the relevant geological features of the selected areas. Continuous and discrete modes have been selected depending on the characteristics and heterogeneity of the sites. For example, the continuous mode has been applied in extended areas in order to have more flexibility and speed to move. In areas with very small-scale heterogeneities the discrete mode has been preferred. In these cases between 5 and 7 measurements have been taken and averaged per point. An advantage of this mode is that it can be measured directly on ground or at a

[Printer-friendly version](#)[Discussion paper](#)

fixed distance above ground. The positions of measuring points and tracks have been georeferenced by a Garmin 62s GPS. The GPS tracks have been used to derive the orientation. For some relatively small mapping areas, like Bahía Glaciares (Site 4; Fig. 1), a grid of 20 x 20 m was previously defined by a tape measure since the error of GPS positions could be in the order of several metres. In these cases, the lines have been used for the orientation. It has to be taken into account that the different data have been obtained from multiple instruments individually without an automatic synchronism. Therefore, all the acquisition units have been manually synchronised, and the sequence of measurement has been done systematically as follows: 1) Marking selected measurement point. 2) GPS measurement with time stamp (with > 6 satellites at direct sight). 3) Removal of the GPS from measurement point to avoid magnetic contamination by this device. 4) Magnetic measurement with time stamp (for G-858). 5) Magnetometer (three axes), accelerometer (three axes), temperature measurement with time stamp (for MOURA). The data files have been pre-processed manually (preliminary corrections of the GPS data with the above-described information). Ad-hoc software has been performed to include the temperature and tilt angles correction of MOURA data, to subtract the Earth geomagnetic field with respect to total intensities in the case of G-858 and each vector components in the case of MOURA, and to plot the different magnitudes of the processed data. Local magnetic field anomalies have been calculated with respect to the International Geomagnetic Reference Field (IGRF) averaged for the month of the surveys. At single sites the surveys were performed during less than 2 hours for which global magnetic field data from a base station and/or from the next magnetic observatories with minute-resolution have been considered for reference (Argentine Islands near Antarctic Peninsula, Port Stanley and Easter Island and Huancayo ~ 1400 km). However, since temporal variations during all survey intervals of single sites were less than ± 10 nT (quiet days), site specific data corrections have not been applied. In all the surveys it has been systematically calculated the correlation parameters between the scalar data of the two magnetometers (G-858 and MOURA) as well as between the vector data of the two separate sensors of MOURA.

[Printer-friendly version](#)[Discussion paper](#)

2.3 Selected sites: mineralogical and geological context The selected test sites for the on ground survey are situated in or near the Southern Andes between latitudes 20°S to 52°S (Fig. 1). Required general site characteristics are that a) exposed rocks are relatively unaltered, b) high resolution grids with scales of metres to centimetres can be performed, and c) exposed rocks are representative for a large number of martian surface rocks.

SITE 1 “El Laco”. Four large magnetite bodies with a total estimated ore resource of 500 million tons crop out around El Laco volcano in the Central Andes (Fig. 2A; Alva-Valdivia et al., 2003; Naranjo et al., 2010). Together with the iron ore deposits of Kiruna (e.g. Jonnsson et al., 2013) they represent worldwide unique examples for very strong local magnetic anomalies, which may be comparable to that observed in some areas of the southern Noachian highlands of Mars (Connerney et al., 2005; Lillis et al., 2013). The selected area with an extension of 0.2 x 0.4 km is situated at the northern margin of the El Laco Sur outcrop (23°50'17"S; 67°29'27" W; 4720 m elevation; Figs. 1 and 2), at the transition between magnetite-bearing ores and early Pleistocene andesitic lava flows which are partly covered by pyroclastic deposits with up to 5 m thickness. The mapping area has not been modified by iron ore mining. Sernageomin (Servicio Nacional de Geología y Minería of Chile; Naranjo et al., 2010) performed aeromagnetic surveys and constructed maps of the anomaly. They reflect a dipolar anomaly according to the isodynamic lines of the map with intensities in the order of ± 150 nT (Alva-Valdivia et al., 2003). However, there is no information available concerning the tracks of these aeromagnetic surveys: spacing and altitudes above ground, because of which, the data are only considered for completeness. Besides, on ground magnetic surveys have not been done previously. Fission-track dating of apatite grown within the magnetites gave an age of 2.1 ± 0.1 Ma (Maksaev et al., 1988). A whole rock age of the andesite from the host rocks of Pico Laco is 2.0 ± 0.3 Ma (Gardeweg and Ramírez, 1985). The origin of the magnetite bodies has been strongly debated. A combined magmatic and hydrothermal origin was proposed by Alva-Valdivia et al. (2003), Sillitoe and Burrows (2002), and Velasco and Tornos (2012). This is based on the fact

[Printer-friendly version](#)[Discussion paper](#)

that field and petrographic evidences suggest that some magnetites have a primarily magmatic texture, whereas others show features which indicate a formation during a hydrothermal triggered re-emplacement of andesitic lava flows. Trace element compositions of the latter magnetite type are not compatible with a magmatic origin. For example, Dare et al. (2014) document that these magnetites are characterized by high Ni/Cr ratios, depleted in Ti, Al, Cr, Zr, Hf and Sc, and show an oscillating zoning of Si, Ca, Mg and rare earth elements. In contrast, oxygen isotope data ($\delta^{18}\text{O}$ of +2 to +4) of many magnetites support a magmatic rather than hydrothermal origin (Jonsson et al., 2013). Microscopy studies under reflected light as well as temperature dependent susceptibility measurements and isothermal remanent magnetization (IRM) acquisition show that low Ti-magnetite and/or maghemite are the magnetic carriers (Alva-Valdivia et al., 2003). Sometimes ilmenite-hematite minerals appear in significant amounts. Grain sizes range from a few microns up to several millimetres. Hysteresis measurements of Alva-Valdivia et al. (2003) of seven ore samples from El Laco Sur point to pseudo-single-domain status and show a large range of Koenigsberger ratios (Q-ratios from 0.02 to > 1000). Paleomagnetic data show distinct local declinations indicating a complex crystallization history, probably during different geomagnetic field orientations (Alva-Valdivia et al., 2003). SITE 2 “Pali Aike”. Lava sheets with volcanic spatter cones represent a common feature in many areas of the surface of Mars (Kereszturi and Németh, 2012; Robbins et al., 2013). On Earth such volcanic rocks often exhibit distinct magnetic anomalies (e.g. Bolos et al., 2012; Urrutia-Fucugauchi et al., 2012). However, only few examples have been mapped with high resolution (e.g. Cassidy and Locke, 2010). Thus, an agglutinated spatter cone of 170 m diameter and surrounding Quaternary lava sheet of the Pali Aike Volcanic Field (PAVF) in southernmost Patagonia (Figs. 1 and 3; Skewes and Stern, 1979) has been selected as potential martian analogue. The well-preserved morphology and stratigraphy indicates an age of approximately 1.0 Ma when considering the succession of various nearby volcanic formations for which ages of 0.16 to 1.5 Ma have been reported (Mejia et al., 2004). The investigated crater is partly filled by pyroclastic material, eolian sediments as well

[Printer-friendly version](#)[Discussion paper](#)

as blocks and detritus, which have been collapsed from the eastern inner crater wall. The mapping site (52°06'43"S; 69°42'28"W; 227 m elevation) covers an area of 400 x 400 m, including the crater and its surroundings (Fig. 3A). SITE 3 "Monturaqui". Impact craters represent a very frequent feature on the Mars surface (Lillis et al., 2013). Depending on e.g. size, target rocks, impactite composition and possible hydrothermal processes they can be characterised by distinct and complex magnetic signatures (e.g. Osinski et al., 2013). On Earth, large impact craters are strongly eroded. In addition, some of them are covered by vegetation or modified by anthropogenic influences. A Late Pleistocene simple type impact crater in the Atacama Desert of northern Chile was selected for this case study (Fig. 1). The crater was discovered in 1962 from aerial photographs and firstly described by Sánchez and Cassidy (1966). It is located at latitude 23°55'40"S and longitude 68°15'42"W at an elevation of 2984 m (Ugalde et al., 2007), has a diameter of 370 m and is 34 m deep (Fig. 4 A, B, C), and was formed during the Quaternary (660 ± 90 kyr BP; Ukstins Peate et al., 2010). Due to the arid climate, it remained morphologically well preserved. The crater has remarkable morphological similarities to the Bonneville impact crater on Mars, which was explored by the Spirit rover of NASA (Grant et al., 2004). Monturaqui target rocks include Jurassic granites cut by some mafic dykes. Both rock types are overlain by a several metre thick sheet of Pliocene ignimbrites. Tiny Fe-Ni-Co-P spherules, all bound in impact glass, have been found within the ejecta blanket. They suggest an iron meteorite as impactor (Bunch and Cassidy, 1972; Kloveranz, 2010). SITE 4 "Bahía Glaciares". Plutonic rocks and layered intrusions form significant parts of the martian crust (e.g. Francis, 2011) and analogues on Earth (McEnroe et al., 2004 and 2009). These rocks may have the capacity to store remanent magnetic signatures that can be used to distinguish between different magmatic rock types during future rover-based magnetic surveys. The Patagonian Batholith in the southernmost Andes provides a good example of continental crust formation on Earth and other planets (Behrmann and Kilian, 2003; Diaz-Michelena and Kilian, 2015). A small mapping area of 20 x 6 m (120 m²) was defined on a Cretaceous granite (Fig. 5; Hervé et al., 2007), which is cross-cut by several mafic

[Printer-friendly version](#)[Discussion paper](#)

North-trending ($\sim 5^\circ\text{N}$) more or less parallel mafic dykes ($52^\circ 48' 28''\text{S}$; $73^\circ 14' 10''\text{W}$; 11 m a.s.l.). This area was chosen because it is a good example of very low intensity magnetic contrast in a small extension, where transition between alternating mafic and felsic outcrops appears at a centimetre-scale.

2.4 Additional analyses

The magnetic field surveys have been complemented with other rock analyses to improve the interpretation of the magnetic signatures of the surveys. Even though the detailed analysis is out of the scope of this work, the types of measurements are briefly described because they support partially some of the conclusions of the work. Firstly, a macroscopic description of the rock types and mineral components has been done at each site. Representative rock samples were collected along the tracks for macroscopic investigation and future analyses in the laboratory. For instance, the samples from Pali Aike (Site 2) and Bahía Glaciares (Site 4) have been analysed with polarization and refracted light microscopy of thin sections of the rocks. Texture and grain sizes of samples from el Laco (Site 1) have been also investigated with a scanning electron microscope (Leo 435 VP, Geology Department, Trier University). The mineral composition of granites and amphibolitic dykes of Bahía Glaciares (Site 4) have been analysed by an X-ray diffractometer (Siemens D500, Geology Department, Trier University). Hysteresis properties of representative samples from Pali Aike (Site 2) and Bahía Glaciares (Site 4) have been characterized magnetically at room temperature by means of a vibrating sample magnetometer at the Space Magnetism Laboratory of INTA, Spain. Magnetic susceptibilities have been also measured with a MS-2 susceptometer by Bartington along the transects at Site 4.

3. Results

The comparative performance and results of the magnetic surveys with MOURA and G-858 magnetometers are described below. In all cases with identical single point measurements the correlation between the two instruments and the two sensors of

[Printer-friendly version](#)[Discussion paper](#)

MOURA has been analysed. The number of differences among the distinct sites has made it possible to demonstrate the versatility of MOURA. Thus, in each study case some of the individual capabilities of the instrument will be highlighted and discussed.

3.1 SITE 1 “El Laco” The surveys were performed with both instruments using continuous and discrete measurement modes. During the surveys the temperature was ranging from 5°C to 27°C. The transects and individual measuring points were georeferenced with the GPS. A Matlab code was used to combine, interpolate and merge the magnetic anomalies and tracks (Fig. 2). The vertical magnetic gradient has been measured at one point where magnetite-bearing ores crop out. Fig. 2B shows an exponential increase of the magnetic anomaly from 1.9 m altitude above the ground down to the rock surface (from + 3,000 to +23,000 nT). The gradient field calculated from the vector components of both vector sensors shows a strong negative vertical component ($m_z = -11.7 \text{ A/m}$) for this point, which has been modelled by a local shallow superficial dipole with a volume of $0.1 \times 0.1 \times 0.4 \text{ m}$, an inclination of -62° and a declination of -67° . The magnetite-bearing outcrops exhibit very high positive magnetic anomalies from 30,000 to $>110,000 \text{ nT}$ while surrounding andesitic lavas and pyroclastic material have much lower positive anomalies (+ 100 to + 2,000 nT). The magnetic anomalies across outcrop transitions between andesites and magnetite-bearing ores have been measured with MOURA in a discrete mode directly on the ground (Fig. 2C) and with a continuous mode (Fig. 2D). The differences in the intensity of andesite anomalies between these two measurements are related to the different and slightly variable distance between the hand-held sensor (25 to 30 cm) and the ground surface of single points (0 cm). The continuous measurements show a large variability of the magnetic anomalies along the ore-bearing outcrops related to either heterogeneous ore compositions or slight variations of the sensor distance from the surface. During G-858 surveys the magnetometer became often saturated when high local magnetic anomalies were reached ($> 80,000 \text{ nT}$). Local surveys with a higher spatial resolution in a continuous mode showed that this situation appeared very frequently and thus did not permit a complete high-resolution survey with this instrument. The surveys

[Printer-friendly version](#)[Discussion paper](#)

with MOURA were not affected by such saturation since this magnetometer has an extended range mode (auto) that doubles the nominal range to $\pm 130,000$ nT per axis, and thus, allows measurements up to higher field intensities as it has been suggested partly for the martian surface. Despite this problem with the G-858 the scalar magnetic maps of both magnetometers are similar with a correlation $R^2 > 0.8$. (comparison in Figs. 2E and 2F). Of relevant importance is that MOURA magnetometer allows the identification of the component that saturates. Since MOURA magnetometer provides vector magnetic data, it is possible to determine the orientation of the field in the area. This is shown in the rosette of Fig. 2G together with the paleodeclinations of other rock samples from El Laco Sur determined by Alva Valdivia et al. (2003).

3.2. SITE 2 “Pali Aike” A dense grid was performed over the depicted surface with G-858 magnetometer (Fig. 3A). In this case, MOURA measurements have been performed with the discrete mode (Fig. 3B) to obtain well-referenced vector data. During the survey, the temperature was oscillating between 7 to 15° C. Fig. 3B compares an interpolated magnetic anomaly map measured with G-858 with discrete points from MOURA that are illustrated with a colour code. Both data sets match very well ($R^2 > 0.82$). A 3-D view of the interpolated magnetic anomalies mapped with G-858 is shown in Fig. 3C. It documents the very high positive anomalies of the crater rim (up to + 12,000 nT). A W-E transect of the crater and its surroundings, and its geological features shows two pronounced positive magnetic anomalies centred at both sides of the crater rim where the agglutinated spatter have been mainly deposited as pillow-like blocks of metre size (Fig. 3D). Vector information obtained with MOURA magnetometer at the individual points along the crater rim and crater infill is illustrated in Fig. 3E. In general the predominant declination in all the measurements taken on consolidated lava blocks and the sedimentary infill is around 355°N (white arrows in Fig. 3E). However, anomalous deviations have been detected in the Eastern and Southern part of the crater (red arrows in Fig. 3E), on single basaltic lava blocks, which have removed into the crater during a post-eruptive collapse of the inner crater wall. MOURA has two magnetometers at a small distance of 10 mm between them. They can be used

[Printer-friendly version](#)[Discussion paper](#)

also to measure some of the components of the gradient of the field. Fig. 3G shows the derivatives respect to z of the components B_x , B_y and B_z of the field. In good agreement with the previous conclusion, the gradient seems to have a homogeneous direction all over the crater with the exception of measurements on single lava blocks that have been removed and re-orientated during the collapse of the wall. For, all these data it has been taken into account the tilt angle of MOURA apart from the deviation respect to the North taken with the GPS. This has been possible due to the fact that MOURA has a tilt angle sensor to measure the deviation from the horizontal. This sensor has been used in this example to derive a gravity contrast along the transect within the crater and along its rim which is illustrated in Fig. 3F. Highest values occur along the eastern and western crater rim whereas lowest values are measured at the western eolian sedimentary crater infill. This relationship is shown in the W-E transect of Fig. 3D.

3.3. SITE 3 “Monturaqui” In this example we performed a dense grid of the centre of the crater as well as the northeastern, eastern and southern part of the crater rim. An interpolated map shows very slight magnetic anomalies (< 50 nT) within the crater (Fig. 4B) whereas more pronounced local negative and positive anomalies from $- 400$ to $> + 600$ nT occur within several meters along the crater rim indicating the existence of metre-sized dipoles. The anomalies are not related to outcrops of exposed granitoids and ignimbrites (Fig. 4C). A higher-resolution mapping was performed at a local area of around 10×20 m at the north-eastern crater rim with both magnetometers in a continuous mode with G 858 and by discrete points with MOURA. Both magnetometers show relatively high positive and negative anomalies ranging from $- 3,500$ up to $> + 6,000$ nT (Fig. 4D). This local field of anomalies indicate the existence of not exposed but near-surface metre-sized dipoles. The anomalies measured with both magnetometers are compared in an X-Y plot of Fig. 4E and indicate a correlation of R^2 of 0.81. The local mapping area at the north-eastern crater rim is characterised by pronounced local topography changes in the range of ± 5 m of elevation. Fig. 4F illustrates that more pronounced negative anomalies occur at topographic lows. This relationship be-

[Printer-friendly version](#)[Discussion paper](#)

tween lower topography points and higher positive anomalies (and vice versa) was also measured with the B1 and B2 sensors of MOURA and is shown in Fig. 4G. This figure documents also the good correlation between both MOURA sensors, B1 showing on average + 1,500 to + 2,000 nT higher values than B2 related to the fact that B1 is 10 mm nearer to the ground surface.

3.4 SITE 4 “Bahía Glaciares” A 20 x 20 m area was mapped with G-858 and MOURA magnetometers along seven high-resolution tracks perpendicular to the dykes (Fig. 5B to 5D) using both continuous and discrete modes. The spacing between the lines is approximately 80 cm and the distance between individual measurement points along the lines range from 5 to 10 cm. All the lines show similar patterns, which allow performing an interpolated map of the area: Figs. 5C and 5D show that the dykes clearly contrast with the granites by slight positive anomalies. Fig. 5E shows one of the high-resolution transects where the magnetic signatures have been measured with both magnetometers and a M2 Bartington device for susceptibilities. The dykes exhibit very clear but weak positive anomalies with respect to the granite in the range from + 20 to + 80 nT. Both magnetometers show similar patterns ($R^2 > 0.8$). Overall slightly higher values (around 30 nT) of MOURA data are attributed to the fact that they have been performed directly on the rock surface whereas those of G-858 magnetometer are measured at a certain distance to the surface (25 to 30 cm). The susceptibility transect shows a very sharp transition at the interfaces between the granite and dykes, the latter having around 70×10^{-6} SI higher values on average. The magnetic anomalies and the susceptibility data show laterally displaced (Fig. 6A, B). This reflects the eastward tilt of the dykes which have dipping angles of 50 to 80° and indicates that the uppermost 2-3 m of the mafic dykes are also integrated within the anomalies measured with both magnetometers while the susceptibility shows only the information of the first centimetres below the surface. Detailed petrographic studies including electron microprobe and XRD analysis show that both granites and amphibolitic dykes do not contain magnetite, but instead include ferromagnetic monoclinic C4 pyrrhotite as magnetic carrier. Areal microscopic mapping of pyrrhotite in thin sections of the different dykes and the

[Printer-friendly version](#)[Discussion paper](#)

granite and XRD analyses of the different rocks indicate a content of 1 to 4 vol.% pyrrhotite with grain sizes ranging from < 5 to $150 \mu\text{m}$. There is a good correlation between the pyrrhotite contents of the different dykes and the amount of the positive anomalies (Fig. 5E).

4.0 Discussion

The results of the different ground magnetic surveys of both magnetometers are discussed with respect to the appropriateness of the different instruments and the relationship to mineralogical and magnetic properties of the exposed rocks. In particular the potential of high-resolution detection of weak magnetic contrast between different surface rock types is considered.

4.1 Instrument Performance Final processed data from both instruments show a very good correlation in intensity of magnetic anomalies (in all cases $R^2 > 0.8$) for the overall measurement range between $-2,000$ nT and $100,000$ nT (Figs. 2E, 2F, 3C, 4E and 5E). The stability of both instruments has been appropriate for the different surveys. G-858 MagMapper shows a better thermal stability, which can be observed during faster temperature variations during the dawn and dusk, when the transducer may experiment thermal variations up to $0.1 \text{ }^\circ\text{C}/\text{min}$. The simultaneous measurement of the temperature and the magnetic field during the prospections diminishes this problem, which can be neglected in areas with magnetic anomalies > 100 nT, but the error can be significant (1 %) in low contrast anomalies (1 nT), also due to the resolution of MOURA instrument. Other ways to compensate the temperature effects could improve these errors (Díaz-Michelena et al., 2015 b). For static measurements the simultaneous temperature measurement corrects very well the magnetic field data (Díaz-Michelena et al., 2015 a). Regarding the dynamic range, both magnetometers have also casted appropriate results in most of the cases. The limitation in this feature affects in a different way the response of both instruments. G-858 is influenced in the measured modulus of the field, while MOURA is affected separately in every axis. Apart from the extension of the range in modulus, this is an advantage since it could provide useful data in two

[Printer-friendly version](#)[Discussion paper](#)

directions despite of saturation in the other axis. For example, at Site 1, the huge intensity of the anomalies makes it impossible to map them with G-858, while MOURA can measure them in the auto mode, when the maximum offset is applied. At the El Laco site MOURA surveys were performed with discrete and continuous modes. The continuous mode enabled a higher resolution and an easier performance but it may include a shifting by slight variations of the distance between sensor and the ground. The extreme high gradient at this site causes a pronounced fluctuation of around $> 5,000$ nT when altitude of the sensor changes from 25 to 30 cm. This can be avoided by discrete measurements directly on the ground, which also enable a better orientation control of the vector sensor. In the case of highly positive anomalies and in combination with high Q-ratios (remanent versus induced magnetic signatures) of the surface rocks, the vector measurements may also have the capability for paleomagnetic implications which is extremely important for planetary exploration. Surface rock alteration processes which modify the magnetic signatures have influence on limited areas. In particular, the related mineral transformations processes are a direct consequence of the contact of the rocks with the hydrosphere and atmosphere, and their influence depth is limited to several tens of metres. This fact together with the exhumation processes offers often the possibility to correlate the measured direction of the magnetization with the coetaneous paleomagnetic field. In the case of Mars, where the main source of field is the remanent magnetization, oriented measurements does not only contain information on the carriers and the possible alteration effects suffered by them, but also record the paleomagnetic field orientation. This possible tool has been applied to remanence dominated sites which is further discussed in section 4.2. The sensor head orientation is also a matter of discussion. Despite the good signal-to-noise ratio presented by the scalar magnetometer, it is affected by the relative orientation of the head and the magnetic field vector. This is highly improved in a 3-axes magnetometer like MOURA. Another consideration is the gradient immunity and capability to derive a gradient of the field of the instruments. On the one hand, MOURA instrument presents a better gradient immunity, which makes it very suitable to map areas with high frequency patching

[Printer-friendly version](#)[Discussion paper](#)

of the signatures. For example, it is very appropriate to perform decimetre-scale resolution mappings like in the cases of El Lago, Monturaqui and Bahía Glaciares. G-858 presents troubles with not so high gradients ($> 20,000$ nT). On the other hand, the inclusion of a second head (and therefore to have two 3-axes magnetometers) in MOURA instrument offers the capability to better understand the characteristics and depths of the sources. This cannot be applied to deep and extended magnetic sources, because the distance between the two magnetometers is very small (10 mm) but it is useful to analyse near surface heterogeneities and will be discussed in section 4.2.

4.2 Capacity for high resolution mapping with tracing of mineralogical and geological characteristics High-resolution ground surveys may indicate compositional variations in soils and/or uppermost crustal rocks, depending on the magnetic contrast between different exposed rocks and the intensity of active magnetic field (Gobashy et al., 2008; Hinze et al., 2013). Despite the fact that it is not the primary goal of MOURA instrument, which is part of the instruments suite of a lander, due to its potential in future exploration missions, the exploration capacity of the commercial G-858 and MOURA magnetometer for extra-terrestrial high-resolution mapping is discussed in the following for the different investigated sites.

El Lago At this site the intensities in the magnetic anomalies range from 0 to + 110,000 nT (Figs. 2C to 2F) which is unique compared to other magnetic mapping results on Earth (e.g. Hinze et al., 2013). The magnetic contrast at the surface transition between andesitic rocks and magnetite-bearing ores is very sharp and extremely high. A change from + 1,200 to + 80,000 nT appears in less than a metre distance. In general, MOURA data show a better definition of the surface rock transitions and local variabilities in the areas with outcrops of magnetite-bearing ores compared to G-858. Furthermore, G-858 could not register some of the anomalies because its response was occasionally saturated (110,000 nT) (Fig. 2D). Some variations in areas with exposed ores (in the order of $\pm 5,000$ nT) could have been caused by slight changes in the sensor distance from the ground during a continuous measuring mode (see Chapter 3.1),

[Printer-friendly version](#)[Discussion paper](#)

the major variations are probably related to the heterogeneous composition and locally distinct magnetic behaviour of the magnetite-bearing ores. The texture of magnetites in the ores indicates in part a primarily volcanic origin, but also frequent recrystallization during later hydrothermal processes can be observed. It is likely that the hydrothermal crystallization took place over a longer period of the Early Quaternary which may have also included magnetic reversals (e.g. Alva-Valdivia et al., 2003; Naranjo et al., 2010). Our field observations and laboratory analyses of collected samples indicate a large scatter in the grain sizes, the porosity content as well as the relative amount of additional apatite (non-magnetic) and pyroxene in these rocks. These features might explain the observed variations. Other variables reported by Alva-Valdivia et al. (2003) include hysteresis parameters and highly variable Q ratios (from 0.01 to >5,000) indicating a wide range of individual properties of the magnetic carriers, compatible with pseudo-single-domain up to multi-domain status. Therefore, we attribute the observed huge variability in magnetic anomalies to local variabilities in the behaviour and magnetic properties of magnetites in near-surface rocks. In areas where andesitic lavas are exposed, field surveys show only low fluctuations of the positive anomalies (Fig. 2C and 2D). This let us to hypothesize that there are no underlying local ore bodies and the lava flows have relatively homogenous compositions. Measurements with the MOURA vector magnetometer show a clear northward declination between 350° to 10° N (Fig. 2G). This value is similar to the present declination of the IRGF at this site (3° N) and can be explained by a very strong induced magnetization consistent with the very high susceptibilities, but relatively low Q ratios (0.01) measured in six of seven samples from El Laco Sur by Alva-Valdivia et al. (2003).

Pali Aike Magnetic surveys have been performed at some volcanoes world-wide with different spatial resolution, e.g. from Australia (Blaikie et al., 2012), New Zealand (Casidy and Locke, 2010) and Italy (Okuma et al., 2009). In general these case studies show more and less positive magnetic anomalies (up to a few thousand nT) depending on the composition of the volcanic rocks, its cooling history and the single versus multi domain status of their magnetites (Clark, 1997). Our example of a small crater (170 m

diameter) and its surroundings at the Pali Aike Volcanic Field was performed with both magnetometers and with a higher spatial resolution than the previous studies. The transect across the Pali Aike crater shown in Fig. 3C and 3D has a spatial resolution of 30-50 cm in-between individual measuring points and thus provides a very good differentiation of different kinds of exposed rocks within the uppermost 1-2 m. Despite the relatively high intensity of the IGRF at Pali Aike (+ 31,000 nT; Fig. 1) the transect across the crater is characterised by very high positive magnetic anomalies of up to + 12,000 nT. These anomalies are mostly pronounced along the crater rim where metre-sized melt spatters have been deposited and cooled down (Fig. 3D). The very strong magnetic signature can be explained by the fact that these lavas contain very frequent tiny magnetite crystals with single domain characteristics in their glassy matrix. The positive anomalies become much lower towards the crater infill and the outer slopes of the crater. An increasing amount of pyroclastic deposits with reorientations during the deposition processes on the steeper slopes of the crater could have reduced the integrated magnetic anomaly of these components. The relatively low local anomalies measured within the crater can be also explained by such multiple re-orientations of magnetic carriers during local redistribution processes including fluvial and eolian activities. Fig. 3E shows arrows for the declination calculated from the vector data of MOURA. The values of all measurements on consolidated lava blocks (white arrows in Fig. 3E) range from 352° to 360° N. These orientations may reflect either the induced present magnetic field or paleofield directions, or a combination of both, depending on their remanence and related Q ratios which reflect often the single versus multidomain status of the basalts. Hysteresis measurements of 25 basaltic lava samples from our mapping area indicate relatively high Q-ratios of 50 to >500 suggesting a strong remanence. Comparable basaltic rocks from other sites worldwide are also characterized by a very strong remanence and a predominant single domain status (Day, 1997; Dos Santos et al., 2015; Dunlop, 2002; Zhao et al., 2006). Our measurements which have been performed directly on detached lava and scoria blocks, clearly collapsed from the inner crater wall, show multiple orientations. They are indicated by red arrows

[Printer-friendly version](#)[Discussion paper](#)

in Fig. 3E and include easterly and westerly declinations. The present field which has a declination of 12°N at Pali Aike, the MOURA field vectors as well as several paleodeclinations from different old Pleistocene lavas of Pali Aike (including magnetic reversals; Mejía et al., 2004) have been compiled in Fig. 3H. The estimated age of the investigated cone is approx. 1.0 Ma which suggests a normal global field for that time. MOURA declinations, ranging from 352° to 360°N , are in a very good agreement with such a normal Bruhns declination as well as other normal declination constrained for other lavas from the Bruhns magnetic period (Mejía et al., 2004) rather than the present field declinations which contrast by $+15^\circ$. This result indicates that MOURA magnetometer could provide paleodeclinations when rocks have a very high remanence (high Q ratios).

Monturaqui impact crater Planetary impact craters can be characterised by a variety of magnetic anomalies which are related in particular to distinct magnetic carriers of the target rocks (e.g. mafic versus felsic or sedimentary) and the sedimentary crater infill as well as the compositions of the impactor, impact-induced melt/glass and/or impact-related hydrothermal mineralization and/or demagnetization (e.g. L'Hereux et al., 2008; Langlais and Thébault, 2011; Osinski and Pierazzo, 2013; Pilkington and Grieve, 1992; Prezzi et al., 2012). At the relatively small Monturaqui crater, a coarse grid of magnetic mapping with spacings of approximately 70 m in-between single measuring points have been previously performed with a caesium magnetometer in the crater and its surroundings by Ugalde et al. (2007). The published interpolated map shows only very low magnetic anomalies in the range of less than ± 200 nT with slightly higher values in the southern and eastern sector of the crater rim. Our grid was measured with a continuous mode of the G-858 and provides a much higher resolution for the central part of the crater as well as its northern, eastern and southern rim and slopes (Fig. 4B). Our data show similar low magnetic anomalies of ± 50 nT for the crater floor as measured by Ugalde et al. (2007), whereas the eastern and north-eastern crater rim is characterised by much stronger local anomalies from -500 to $> + 600$ nT. The transitions of the outcrops of granitic and ignimbritic target rocks shown Fig. 4C can-

[Printer-friendly version](#)[Discussion paper](#)

not be depicted by the distinct anomalies, while two northwest to southeast-trending dykes seems to responsible for some anomalies at the eastern crater rim (Fig. 4B). In general, the crater rim is characterised by a patchwork of pronounced local positive and negative anomalies which can be caused by small-scale near-surface dipoles which are probably not exposed. Not exposed fragments of the impactor, for which a diameter of ~ 15 m has been modelled (Echaurren et al., 2005), represent a potential source for these magnetic anomalies. Fe-Ni spherules which have been found in impact melt/glass fragments along the eastern and western crater rim has been classified as a Group I Coarse Octahedrite (Bunch and Cassidy, 1972; Buchwald, 1975) and contain schreibersite, cohenite, fossil taenite, and kamacite as major components as well as some troilite. Laboratory analyses of these components show both high remanent magnetization as well as very high susceptibility (Ugalde et al., 2007; Ukstins Peate et al., 2010). The very pronounced negative to positive anomalies from - 2,000 to + 5,000 nT which have been mapped in a local area of 50 to 100 m extension at the northeastern crater rim (Fig. 4D) also require near-surface rocks with strong dipoles. Not exposed metre-sized fragments of the iron-bearing impactor represent the most likely explanation. In this local area topographic highs are formed by ignimbrite blocks of low-magnetic signature which has been ejected from the crater and deposited on the rim during the impact event. Topographic highs formed by these blocks cause a higher distance of the magnetic sensor with respect to the probably underlying fragments of the impactor while measurements in topographic lows show much higher positive anomalies (Fig. 4F and 4G).

Bahía Glaciares Mafic dykes within felsic to intermediate crustal rocks often produce pronounced local positive magnetic anomalies since they include frequent tiny magnetites (e.g. Hinze et al., 2013). However, in our case study the petrographical investigations indicate that the investigated dykes have not preserved their original magmatic mineral textures and the mineral assemblage point to an emplacement and later equilibration of the dykes under upper greenschist to amphibolite facies conditions (Bucher and Grapes, 2011; Philpotts and Ague, 2009). Granites and dykes do not contain

magnetite, but both contain monoclinic pyrrhotite as magnetic carrier (Dekkers, 1988, 1989; Clark, 1984). This mineral appears disseminated and along veins and has been formed during hydrothermal mineralization together with Cu and Au enrichments during the exhumation (Díaz-Michelena and Kilian 2015; Nelson, 1996; Schalamuk et al., 1997). Despite the lower potential of pyrrhotite to produced magnetic anomalies both magnetometers (MOURA and G-858) clearly show high resolution and slightly positive magnetic anomalies (+ 30 to + 80 nT) as well as higher susceptibilities of the dykes compared to the granites (Fig. 5E). The surface transitions between dykes and granites are characterised by very sharp anomalies within a decimetre scale. The amount of pyrrhotite (1 to 4 vol. %) which has been quantified in samples of the granite and seven dykes shows a very good correlation with the intensity of the positive anomaly (from + 160 to + 220 nT; Fig. 5F). This result confirms the potential of both magnetometers to explore local mineral enrichments which are often produced by hydrothermal processes associated with gold and copper enrichments (Direen et al., 2008). Hinze et al. (2013) show examples of mafic dykes in felsic rocks where different dyke geometries cause distinct shapes of local magnetic anomalies across dykes. For the investigated Bahía Glaciares site Fig. 6 illustrates asymmetric behaviour of the magnetic anomalies along lines which have been measured across dykes which dip with between 50 to 80°. All anomalies are slightly displaced towards the shallower dipping site of the dykes indicating an integration of the magnetic signature of the uppermost 2-3 m of not exposed dykes. This fact together with the difference in contrast obtained with the magnetometers and the susceptometer (Fig. 5) points out that the magnetic field measurements average large volume sources and can lead to wrong conclusions if used as a quantitative mineralogical marker. These results indicate that a multihead (magnetometer and susceptometer) instruments would provide much better results for this purpose.

5. Conclusions

Several sites with a huge variability in magnetic anomalies have been analysed. As a

[Printer-friendly version](#)[Discussion paper](#)

first conclusion it can be said that the surface measurement of the sourced field often gives direct information on the composition, petrogenesis and alteration processes of the surface rocks. For the study, two different magnetometers have been used. On the one hand MOURA vector magnetometer and gradiometer (< 200 g: 72 g instrument + batteries and control PC), developed for Mars surface measurements, as the demonstration of the technology for planetary surveys, and on the other hand a commercial caesium G-858 magnetometer (8 - 9 kg), also used as a reference. The studied areas are considered Mars analogues and they are representative of the intensity range of the expected anomalies on the Red Planet crust. According to the comparison with the reference instrument, it has been demonstrated that MOURA magnetometer is not only appropriate for the static measurement of the absolute value of the magnetic field and its temporal variations, but is also suitable for the prospection measurements in the range of sourced fields from < 15,000 to > 120,000 nT and to reproduce the magnetic contrast of the terrains. Furthermore, MOURA offers vector data of the field and components of the gradient with a significantly lower mass. The particular conclusions for the four case studies are the following: 1) El Laco magnetite-bearing ore deposits in the Northern Andes of Chile represents a world-wide unique example with extremely high on ground anomalies ranging from + 30,000 to + 110,000 nT, which may be comparable to highly magnetic rocks of the Noachian martian crust. In this case MOURA enabled better results than G-858 due its larger range of operation (130,000 nT /axis). The declinations measured also by MOURA vectors represent the active global field due to induced-dominated magnetic rock properties with very low Q-ratios. 2) A crater in the Pali Aike Volcanic Field, in southern Chile, shows very high positive magnetic anomalies (up to 12,000 nT) of its crater rim caused by frequent tiny and single domain magnetite crystals in the matrix of basaltic lava spatters. Since these rocks, like that of many other comparable volcanic rocks on Earth and other planets, have high Koenigsberger ratios (Q) and thus are dominated by their remanence, MOURA vector data have been used to determine the paleomagnetic orientation during the crater formation around 1 Ma before present. In addition, the different later reorientations

[Printer-friendly version](#)[Discussion paper](#)

of single lava blocks during their collapse from the steep inner crater wall could have been constrained by the vector data. 3) The small Monturaqui impact crater in the Atacama Desert of Northern Chile represents an analogue for many other simple type craters, like Bonneville crater on Mars. The granitoid and rhyolitic target rocks have few magnetic carriers and only weak magnetic anomalies. Pronounced anomalies along the crater rim indicate metre-sized unexposed remnants of the iron-bearing impactor (octahedrite). Local mapping with a decimetre resolution, with intensities ranging from $-2,000$ to $+6,000$ nT, corroborates the existence of such localised, strong and relatively small size (1 metre) dipoles (iron meteorite fragments) near the surface. 4) A site within the Patagonian Batholith of the southernmost Andes provides a window into deeper planetary crustal magnetic signatures. The exposed rocks include granites and mafic dykes that have been partly equilibrated at lower amphibolite facies conditions, where all primary magmatic magnetites have been transformed to iron-bearing silicates, and a later hydrothermal mineralization produced pyrrhotite as only magnetic carrier. The freshly exposed transitions between these granites and mafic dykes have been mapped on a decimetre-scale. Despite the very low magnetic contrast from 20 to 80 nT both rock types could have been clearly distinguished. In addition, the amount of pyrrhotite, ranging from 1 to 4 vol. %, is well correlated with the positive magnetic anomalies of the dykes. This documents the potential for mapping of hydrothermal mineralization processes as well as associated gold and copper enrichments, even if the magnetic contrast is very low.

Acknowledgements: Authors acknowledge all MOURA MetNet team; in particular, the payloads electronics engineering laboratory for their work with the magnetometer and V. Apéstigue for the technical support. This work was supported by the Spanish National Space Programme (DGI-MEC) through the project AYA2011-29967-C05-01 and the Spanish National Space Program of R&D Externalization through the project PRI-PIBUS-2011-1105.

References: Acevedo R., Rabassa M., Ponce J., Martínez O., Orgeira M., Prezzi

[Printer-friendly version](#)[Discussion paper](#)

C., Corbella H., González-Guillot M., Rocca M., Subías I. and Vásquez C.: The Bajada del Diablo astrobleme-strewn field, Argentine Patagonia: extending the exploration to surrounding areas, *Geomorphology*, 169-170, 151-164, 2012. Acuña M. H., Connerney J. E. P., Wasilewski P., Lin R. P., Anderson K. A., Carlson C. W., McFadden J., Curtis D. W., Mitchell D., Reme H., Mazelle C., Sauvaud J. A., d'Uston C., Cros A., Medale J. L., Bauer S. J., Cloutier P., Mayhew M., Winterhalter D., Ness N. F.: Magnetic Field and Plasma Observations at Mars: Initial Results of the Mars Global Surveyor Mission, *Science*, 279(5357), 1676-1680, 1998. Alva-Valdivia, L.M., Rivas, M.L., Goguitchaichvili, A., Urrutia-Fucugauchi, J., González, J-A., Morales, I, Gómez, S., Henríquez, F., Nyström, J.O. and Naslund, H.R.: Rock-magnetic and oxide microscopic studies of the El Laco iron-ore deposits, Chilean Andes, and implications for magnetic anomaly modelling, *International Geology Review*, 45 (6), 533 – 547, 2003. Balta, J.B. and McSween, H.Y.: Water and the composition of Martian magmas, *Geology*, 41(10), 1115 - 1118, 2013. Blaikie, T.N., Ailleres, L., Cas, R.A.F. and Betts, P.G.: Three-dimensional potential field modelling of a multi-vent maar-diatreme at The Lake Coragulac maar, Newer Volcanics Province, south-eastern Australia, *Journal of Volcanology and Geothermal Research*, 235–236, 70–83, 2012. Bolós, X., Barde-Cabusson, S., Pedrazzi, D., Martí, J., Casas, A., Himi, M. and Lovera, R.: Investigation of the inner structure of La Crosa de Sant Dalmai maar (Catalan Volcanic Zone, Spain), *Journal of Volcanology and Geothermal Research*, 247–248, 37–48, 2012. Breitsprecher, K. and Thorkelson, D.J.: Neogene kinematic history of Nazca–Antarctic–Phoenix slab windows beneath Patagonia and the Antarctic Peninsula, *Tectonophysics*, 464, 10 – 20, 2009. Breuer, S., Kilian, R., Weinrebe, W., Schörner, D. and Behrmann, J.: Glacier and tectonic control on fjord morphology and sediment deposition in the Magellan region (53°S), *Marine Geology*, 346, 31 – 46, 2013. Bucher, K., and Grapes, R.: *Petrogenesis of metamorphic rocks*. Springer, 8th edition, 428 pp., Berlin-Heidelberg, DOI 10.1007/978-3-540-74169-5, 2011. Buchwald, W.: *Handbook of Iron Meteorites, Their History, Distribution, Composition, and Structure, Volume 3*. Berkeley, University of California Press, 1975. Bunch, T.E. and Cassidy, W.A.: *Petro-*

[Printer-friendly version](#)[Discussion paper](#)

graphic and electron microprobe study of the Monturaqui Impactite, *Contrib. Mineral. Petrol.* 36, 95-112, 1972. Carter, J. and Poulet, F.: Ancient plutonic processes on Mars inferred from the detection of possible anorthositic terrains, *Nature Geoscience*, 6, 1008 – 1012, 2013. Cassidy, J. and Locke, C.A.: The Auckland volcanic field, New Zealand: Geophysical evidence for structural and spatio-temporal relationships, *Journal of Volcanology and Geothermal Research*, 195, 127-137, 2010. Clark, D.A.: Hysteresis properties of sized dispersed monoclinic pyrrhotite grains, *Geophysical Research Letters*, 11, 173 – 176, 1984. Clark, D.A.: Magnetic petrophysics and magnetic petrography: aids to geological interpretation of magnetic surveys, *AGSO Journal of Australian Geology and Geophysics*, 17, 83 – 103, 1997. Connerney, J.E.P., Acuña, M.H., Ness, N.F., Kletetschka, G., Mitchell, D.L., Lin, R.L. and Reme, H.: Mars magnetic field - Results from the Mars Global Surveyor, *PNAS*, 102 (42), 14970-14975, 2005. Dare, S.A.S., Barnes, S.-J. and Beaudoin, G.: Did massive magnetite “lava flows” of El Laco (Chile) form by magmatic or hydrothermal processes? New constraints from magnetite composition by LA-ICP-MS, *Mineral Deposita*, Doi 10.1007/s00126-014-0560-1, 2014. Dekkers, M.J.: Magnetic properties of natural pyrrhotite. Part I: behaviour of initial susceptibility and saturation-magnetization-related rock-magnetic parameters in a grain size dependent framework, *Physics of the Earth and Planetary Interiors*, 52, 376 – 393, 1988. Dekkers, M.J.: Magnetic properties of natural pyrrhotite. Part II: high and low temperature behaviour of Jrs and TRM as function of grain size, *Physics of the Earth and Planetary Interiors* 57, 266 – 283, 1989. Díaz-Michelena, M., Sanz, R., Cerdán, M. F. and Fernández, A.B., 2015. Calibration of QM-MOURA three-axis magnetometer and gradiometer, *Geosciences Instrumentation, Methods and Data Systems*, 4, 1-18, 2015a. Díaz-Michelena, M. and Kilian, R.: MOURA Martian magnetometer potential for high resolution magnetic mapping, *General Assembly 2013. Geophysical Research Abstracts*, Vol. 15, EGU2013-1886, 2013. Díaz Michelena, M., Cobos Arribas, P. and Aroca Hernández-Ros, C. Lock-in amplifiers for AMR sensors, *Sensors and Actuators A: Physical* 222, 149-159, 2015b. Díaz-Michelena, M. and Kilian, R.: Magnetic signatures of the orogenic crust of the

[Printer-friendly version](#)[Discussion paper](#)

Patagonian Andes with implication for planetary exploration, *Physics of the Earth and Planetary Interiors*, doi:10.1016/j.pepi.2015.08.005, 2015. Dieren, N.G., Pfeiffer, K.A. and Schmidt, P.W.: Strong remanent magnetization in pyrrhotite: A structurally controlled example from the Paleoproterozoic Tanmi orogenic gold province, northern Australia, *Precambrian Research* 165, 96 – 106, 2008. Dos Santos, E., Gattacceca, J., Rochette, P., Scorzelli, R.B., Fillion, G. et al. Magnetic hysteresis properties and 57Fe Mössbauer spectroscopy of iron and stony-iron meteorites: Implications for mineralogy and thermal history, *Physics of the Earth and Planetary Interiors*, 242, 50-64, 2015. Dunlop, D.J.: Theory and application of the Day plot (Mrs/Ms versus Hcr/Hc). 2. Application to data for rocks, sediments, and soils, *Journal of Geophysical Research*, 107 (B3), 10.1029/201JB000487, 2002b. Echaurren, A., Ocampo, C., Rocca, M.C.L.: A mathematic model for the Monturaqui impact crater, Chile, South America, 68th Annual Meteoritical Society Meeting, 5004 p., 2005. Eppelbaum, L.V.: Quantitative Interpretation of Magnetic Anomalies from Bodies Approximated by Thick Bed Models in Complex Environments, *Environmental Earth Sciences*, 74, (7), 5971-5988, 2015. Eppelbaum L.V. and Mishne A.R: Unmanned airborne magnetic and VLF investigations: effective geophysical methodology of the near future, *Positioning*, 2(3), 112–133, 2011. Francis, D.: Columbia Hills - An exhumed layered igneous intrusion on Mars? *Earth and Planetary Science Letters*, 310, 59 – 64, 2011. Gardeweg, M., and Ramírez, C. F.: Hoja Rio Zapaleri, II Region de Antofagasta: Santiago, Chile, Servicio Nacional de Geología y Minería, Carta Geológica de Chile, 66, 89 p., 1985. Gobashy, M.M. and Al-Garni, M.A.: High Resolution Ground Magnetic Survey (HRGM) for determining the optimum location of subsurface dam in Wadi Nu'man, Makkah Al Mukarramah, KSA, *JKAU: Earth Sci.*, 19, 57-83, 2008. Grant, A. J., Arvidson, R., Bell III, J.F., Cabrol, N.A., Carr, M.H., Christensen, P., Crumpler, L., Des Marais, D. J., Ehlmann, B. L., Farmer, J., Golombek, M., Grant, F.D., Greeley, R., Herkenhoff, K., Li, R., McSween, H. Y., Ming, D. W., Moersch, J., Rice Jr., J. W., Ruff, S., Richter, L., Squyres, S., Sullivan, R. and Weitz, C. : Surficial Deposits at Gusev Crater Along Spirit Rover Traverses, *Science*, 305 (5685), 807–810, 2004. Greenough, J.D. and

[Printer-friendly version](#)[Discussion paper](#)

Yacoby, A.: A comparative geochemical study of Mars and Earth basalt petrogenesis, *Can. J. Earth Sci.*, 50, 78-93, 2013. Groschner, C., Lan, S., Wise, A., Leary, A., Lucas, M., Park, C., Laughlin, D., Díaz-Michelena, M. and McHenry, M.: The role of atmosphere on exsolution processes and magnetic properties of ulvospinel, *IEEE Transactions on Magnetism* 49(7), 4273 – 4276, 2013. Hervé, F., Pankhurst, R.J., Fanning, C.M., Calderón, M. and Yaxley, G.M.: The South Patagonian batholith: 150 my of granite magmatism on a plate margin, *Lithos*, 97, 373 – 394, 2007. Hinze, W.J., Von Frese R.R.B. and Saad, A.H.: Gravity and magnetic Exploration, Principles, Practices and Applications, Cambridge University Press, 525 p., 2013. Ialongo, S., Fedi, M. and Florio, G.: Invariant models in the inversion of gravity and magnetic fields and their derivatives. *Applied Geophysics*, 110, 51–62, 2014. Jonnsson, E., Troll, V.R., Högdahl, K., Harris, C., Weis, F., Nilsson, K. and Skelton, A. Magmatic origin of giant 'Kiruna-type' apatite-iron-oxide ores in Central Sweden, *Scientific Reports*, 3, 1644, [Doi:10.1038/srep01644](https://doi.org/10.1038/srep01644), 2013. Kereszturi, G. and Németh, K.: Monogenetic Basaltic Volcanoes: Genetic Classification, Growth, Geomorphology and Degradation, In: Németh, K. (ed.): *Updates in Volcanology - New Advances in Understanding Volcanic Systems*, ISBN: 978-953-51-0915-0, InTech: 3-88. DOI: 10.5772/51387, 2012. Kilian, R. and Behrmann, J. H.: Geochemical constraints on the sources of continent-related deep sea sediments and their recycling in arc magmas of the Southern Andes, *Journal of Geol. Soc. London*, 160, 57-70, 2003. Klobberdanz, C. M.: Geochemical analysis of the Monturaqui Impact Crater, Chile, Ph thesis, University of Iowa, 2010. Lan, S., Groschner, C., Runco, J., Wise, A., Díaz-Michelena, M., Laughlin, D. and McHenry, M.: Phase identification and temperature-dependent magnetization of Ti-rich titanomagnetite in different atmospheres magnetism, *IEEE Transactions on Magnetism*, 49 (7), 4314 – 4318, 2013. Langlais B. and E. Thébaud, E.: Predicted magnetic signatures of Martian (de)magnetized impact craters and a statistical comparison with observations, *Icarus*, 212(2), 568-578, 2011. L'Heureux, E., Ugalde, H., Milkereit, B., Boyce, J., Morris, W., Eyles, N., and Artemieva, N.: Using vertical dikes as a new approach to constraining the size of buried craters: An

[Printer-friendly version](#)[Discussion paper](#)

example from Lake Wanapitei, Canada, in Kenkmann, T., Hörz, F., and Deutsch, A., eds., Large meteorite impacts III: Geological Society of America Special Paper 384, p. 43 – 50, 2005. Lillis, R.J., Robbins, S., Manga, M., Halekas, J.S. and Frey, H.V.: Time history of the Martian dynamo from crater magnetic field analysis, *Journal Geophysical Research: Planets*, 118, 1–24, doi:10.1002/jgre.20105, 2013. MaksaeV, V., Gardeweg, M., Ramírez, C. F. and Zentilli, M.: Aplicación del método trazas de fisión (fission track) a la datación de cuerpos de magnetita de El Laco e Incahuasi en el altiplano de la región de Antofagasta, Congreso Geológico Chileno 5th, Santiago, v. 1, p. B1–B23, 1988. McEnroe, S.A., Brown, L.L. and Robinson, P. (2004): Earth analog for Martian magnetic anomalies: remanence properties of hemo-ilmenite norites in the Bjerkreim-Sokndalintrusion, Rogaland, Norway, *Journal of Applied Geophysics* 56: 195– 212. McEnroe S.A., Fabian, K., Robinson, P., Gaina, C. and Brown, L.L.: Crustal magnetism, lamellar magnetism and rocks that remember, *Elements*, 5, 241 – 246, 2009. Mejia, V., Opdyke, N.D., Vilas, J.F., Singer, B.S. and Stoner, J.S.: Plio-Pleistocene time-averaged field in southern Patagonia recorded in lava flows, *Geochemistry Geophysics Geosystems*, 5(3), DOI:10.1029/2003GC000633a, 2004. Morschhauser, A., Lesur, V. and Grott, M. A spherical harmonic model of the lithospheric magnetic field of Mars, *Journal of Geophysical Research*, 119, 6, 1162-1188, 2014. Naranjo, J.A., Henríquez, F. and Nyström J.O.: Subvolcanic contact metasomatism at El Laco Volcanic Complex, Central Andes, *Andean Geology*, 37, 110-120, 2010. Nelson, E.P.: Suprasubduction Mn mineralization: Metallo-tectonic Terranes of the Southernmost Andes, pp. 315-330, In *Geophysical Monograph* 98, Eds.: Bebout, G.E. et al., 1996. Okuma, S., Stotter, C., Supper, R., Nakatsuka, T., Furukawa, R. and Motschka, K.: Aeromagnetic constraints on the subsurface structure of Stromboli Volcano, Aeolian Islands, Italy, *Tectonophysics*, 478, 19–33, 2009. Osinski, G.R. and Pierazzo, E.: *Impact cratering: Processes and products*, Wiley, 330 p., 2013. Philpotts, A. and Ague, J.: *Principles of Igneous and Metamorphic Petrology*, 684 p., Cambridge University Press, 2009. Pilkington, M. and Grieve, R.: The geophysical signature of terrestrial impact craters, *Reviews of Geophysics*, 30, 161-181, 1992. Robbins, S.J.,

[Printer-friendly version](#)[Discussion paper](#)

Hynek, B.M., Lillis, R.J. and Bottke, W.F.: Large impact crater histories of Mars: The effect of different model crater age techniques, *Icarus*, 225, 173–184, 2013. Prezzi C., Orgeira M., Acevedo R., Ponce J., Martinez O., Rabassa J., Corbella H., Vásquez C., González-Guillot M. and Subías I.: Geophysical characterization of two circular structures of Bajada del Diablo (Argentina): indication of impact origin, *Physics of the Earth and Planetary Interiors*, 192-193, 21-34, 2012. Sánchez, I. and Cassidy, W.: A previously undescribed meteorite crater in Chile, *Journal of Geophysical Research*, 71, 4891-4895, 1966. Schalamuk, I.B., Zubia, M., Genini, A. and Fernández, R.R.: Jurassic epithermal Au-Ag deposits of Patagonia, Argentina, *Ore Geology Reviews* 12, 173 – 186, 1997. Sillitoe, R.H. and Burrows, D.R.: New field evidence bearing on the origin of the El Laco magnetite deposits, Northern Chile, *Economic Geology*, 97, 1101-1109, 2002. Skewes, M.A. and Stern C.R.: Petrology and geochemistry of alkali basalts and ultramafic inclusions from the Pali-Aike Volcanic Field in southern Chile and the origin of the Patagonian Plateau lavas, *Journal of Volcanology and Geothermal Research*, 6, 3-25, 1979. Stolpe, E.M., Baker, M. B., Newcombe, M. E., Schmidt, M. E., Treiman, A. H., Cousin, A., Dyar, M. D., Fisk, M. R., Gellert, R., King, P. L., Leshin, L., Maurice, S., McLennan, S. M., Minitti, M. E., Perrett, G., Rowland, S., Sautter, V., Wiens, R. C. , MSL Science Team: The petrochemistry of Jake_M: A Martian Mugearite, *Science*, 341, 6153. DOI: 10.1126/science.1239463, 2013. Ugalde, H., Valenzuela, M. and Milkereit, B.: An integrated geophysical and geological study of the Monturaqui impact crater, Chile, *Meteoritics & Planetary Science*, 42(12), 2153-2163, 2007. Ukstins Peate, I., Klobberdanz, C., Peate, D.W., Chung Wan, L., Cabrol, N., Grin, E., Piatek, J. and Chong, G.: Non-modal melting of target rocks to produce >ite at Monturaqui Crater, Chile, 41st Lunar and Planetary Science Conference, 1-5 March 2010, Houston, Texas, 2010. Urrutia-Fucugauchi, J., Trigo-Huesca, A. and Pérez-Cruz L.: Magnetic links among lava flows, tuffs and the underground plumbing system in a monogenetic volcano, derived from magnetics and paleomagnetic studies, *Physics of the Earth and Planetary Interior*, 212-213, 10-18, 2012. Valesco, F. and Tornos, F.: Insights on the effect of hydrothermal alteration in the El Laco magnetite deposits,

[Printer-friendly version](#)[Discussion paper](#)

Revista de la Sociedad Espanola de Mineralogía, 16, 2012. Wray, J.J., Hansen, S.T., Dufek, J., Swayze, G.A., Murchie, S.L., Seelos, F.P., Skok, J.R., Irwin III, R.P. and Ghiorso, M.S.: Prolonged magmatic activity on Mars inferred from the detection of felsic rocks, Nature Geoscience, 6, 1013 – 1017, 2013. Zhao, Xixi, et al. Unraveling the magnetic carriers of igneous cores from the Atlantic, Pacofoc, and the southern Indian oceans with rock magnetic characterization, Physics of the Earth and Planetary Interiors, 156, 294-328.

Please also note the supplement to this comment:

<http://www.geosci-instrum-method-data-syst-discuss.net/gi-2015-25/gi-2015-25-AC1-supplement.pdf>

Interactive comment on Geosci. Instrum. Method. Data Syst. Discuss., doi:10.5194/gi-2015-25, 2016.

[Printer-friendly version](#)

[Discussion paper](#)

