



1	Soil Salinity Mapping and Hydrological Drought Indices Assessment in Arid Environments
2	<b>Based on Remote Sensing Techniques</b>
3	Mohamed Elhag and Jarbou A. Bahrawi
4	Department of Hydrology and Water Resources Management, Faculty of Meteorology,
5	Environment & Arid Land Agriculture, King Abdulaziz University
6	Jeddah, 21589. Saudi Arabia.
7	Correspondence to: <u>melhag@kau.edu.sa</u>

8

### 9 Abstract

Vegetation indices are mostly described as crop water derivatives. Normalized Difference 10 Vegetation Index (NDVI) is one of the oldest remote sensing applications that we ly used to 11 evaluate crop vigor directly and crop water relationships indirectly. Recently, several NDVI 12 derivatives are exclusively used to assess crop water relationships. Four hydrological drought 13 indices are examined in the current research study. Water Supply Vegetation Index (WSVI), Soil 14 15 Adjusted Vegetation Index (SAVI), Moisture Stress Index (MSI) and Normalized Difference Infrared Index (NDII) are implemented in the current study as an indirect tool to map the effect 16 of different soil salinity level on crop water stress in arid environments. In arid environments? 17 18 such as Saudi Arabia, water resources are under pressure especially groundwater levels. Groundwater wells are rapidly depleted due to the heavy abstraction of the reserved water. 19 Heavy abstractions of groundwater, which exceed crop water requirements in most of the cases 20 21 are powered by high evaporation rates in the designated study area because of the long days of 22 extremely hot summer. Landsat OLI-8 data were extensively used in the current research to obtain several vegetation indices in response to soil salinity in Wadi Ad-Waser. Principal 23 Component Analysis and Artificial Neural Network Analysis are complementary tools to 24 25 understand the regression pattern of the hydrological drought indices in the designated study 26 area.

27

Keywords: Arid Environment, Remote Sensing techniques, Soil Salinity Mapping, VegetationIndices.





#### 30 **1. Introduction**

Remote sensing data sidered to be a convenient source to perfume several vegetation indices in either simple or complicated band ratio combinations. Satellite images offer a large amount of data that could be analyzed, processed and stored to better understand several vegetation indices based on the type of the satellite sensor used (Govaerts et al., 1999; Pinty et al., 2009). Hypothetical backgrounds have been implemented to improve and enhance the optimization of particular satellite sensor to support certain vegetation indices (Verstraete et al., 1996; Gobron et al., 2000; Psilovikos and Elhag, 2013).

Spectral vegetation indices are mathematical combinations of different spectral bands mostly in the visible and near-infrared regions of the electromagnetic spectrum. Vegetation activities can be measured comprehensively through semi-analytical methods of spectral band ratios that have been extensively used to detect not only seasonal variability of the vegetation cover but also local scale spatial variability (Broge and Mortensen, 2002; Xiao et al., 2002).

The generic principle of utilizing vegetation indices is to improve the interpretation of the spectral data reflected from a vegetation cover. Spectral reflectance variabilities tend to differentiate between different vegetation characteristics based on crop water relationships and other surrounding features of soil components and atmosphere based on the maximization of vegetation characteristics over the surrounding environments (Moulin and Guerif, 1999; Boegh et al., 2002). Color, roughness, and water content are mainly the soil components that affect soil spectral reflectance (Curran, 1983a, b; Bouman and Tuong, 2001).

50 Soil parameter variation tends to draw a line on a plenary scattergram. Nevertheless, this line 51 used as a reference point and known as "soil line" in vegetation studies involved both Red and 52 Infrared spectral bands (Colombo et al., 2003, Elhag, 2014a, b). Utilization of vegetation indices





has been challenged always by the najor difficulty which is the minimization of soil component 53 interferences and sensitivity maximization of atmospheric variations (Leprieur et al., 1994; Oi et 54 al., 1994). Atmospherically Resistant Vegetation Index (ARVI) developed by Kaufman and 55 Tanré (1992) and the Global Environmental Monitoring Index (GEMI) developed by Pinty and 56 Verstraete (1991) are the less sensitive vegetation indices to the atmospheric variation. On the 57 other hand, Qi et al. (1994) reported that the (GEMIT) is soil noise sensitive. Higher noise 58 sensitivity of GEMI has completely disabled the index and classifies it to arid region 59 Implementations of vegetation indices are varied from a local leaf scale to continental vegetation 60 scale. Moreover, certain indices tend to be site and/or species specific (Clevers, 1989; Elhag 61 2014a) and it can't be applied not only to different species but also different leave structure and 62 canopies geometry (Xiao et al., 2002). Scholarly work of Kerr, and Ostrovsky (2003), Pettorelli 63 et al. (2005), Huete et al. (2008) and Elhag (2014b) reported that several vegetation indices detailed and the several vegetation indices detailed at the 64 to estimate different vegetation parameters extensively includes: Leaf Area Index (LAI), 65 Fractional Vegetation Cover (FC), Crop Water Shortage Index (CWSI), Drought Severity Index 66 (DSI) and Water Supply Vegetation Index (WSVI). 67

Soil salinization is a dynamic process design basically when an excess of irrigational water is 68 frequently used in the drainage capacity of the fields (Wardlow and Egbert, 2010). 69 Implementations of remote sensing techniques in soil salinity mapping achieved comprehensive 70 71 results on the regional scale (Montandon and Small, 2008). Brightness Index (BI), Normalized Difference Salinity Index (NDSI) and Salinity Index (SI), are widely distinguished in soil salinity 72 mapping in an arid environment (Douaoui et al., 2006; Jiapaer et al., 2011). Frent research 73 74 aims to evaluate the suitability of different vegetation indices for a different level of remotely sensed soil salinity with contrasting to crop water relationships in Wadi Ad-Wasser. 75







### 76 2. Materials and methods

#### 77 **2.1. Study area**

The study area, Wadi Ad Dawasir town ocated in the plateau of Najd at Lat 440 43' and Lon 78 200 29'; about 300 km south of the capital city Riyadh. The study area illustrated in Figure 1 is 79 comprised of gravelly tableland disconnected by insignificant sandy oases and isolated mountain 80 bundles. Across the Arabian Peninsula whole, the tableland slopes toward the east from an 81 82 elevation of 1,360 meters in the west to 750 meters at its easternmost limit. Wadi Ad Dawasir and Wadi ar Rummah most important pattern of the ancient riverbeds remains in the study 83 area. Wadi Ad Dawasir and Najran regions are the major irrigation water abstruction from Al-84 Wajid aquifer. Agriculture in Wadi Ad Dawasir area consists of technically highly developed 85 farm enterprises that operate dern pivot irrigation system. The size of center pivot ranges 86 from 30 ha to 60 ha with farms managing hundreds of them with the corresponding number of 87 wells. The main crop grown in winter is wheat and occasionally potatoes, tomatoes or melons. 88 All year fodder consists of alfalfa, which is cut up to 10 times a year for food. Typical summer 89 crops for fodder are sorghum and Rhodes grass, which is perennial, but dormant in winter. The 90 91 shallow alluvial aquifers could not sustain the high groundwater abstraction rates for a long time 92 and groundwater level declined dramatically in most areas. Meteorological features of the area are speckled. Five elements of meteorology are constantly recorded through fixed weather 93 94 station located within the study area. Temperature varies from 6° C as minimum temperature to 43° C as maximum temperature. Relative humidity is mostly stable at 24 %. Solar radiation of 95 average sunrise duration is generally 11 hereity. Average wind speed is closer to 13 km/h and 96 may reach up to 46 km/hr in thunderstorm incidents. Finally, mean annual rainfall is about 37.6 97 mm (Al-Zahrani and Baig, 2011). 98





99

### Figure 1. Location of the study area (Elhag, 2016).

100	2.2.	Meth	odolo	gical	framewor	rk
			0	B		

- 101 The current research work is based on assessing a regression correlation between different
- 102 vegetation indices and their spatial corresponding soil salinity values conducted from satellite
- 103 images. Principal Component Analysis is the undertaken tool to envisage the impacts of Soil
- 104 Salinity on the current vegetation.

### 105 **2.3. Estimation of vegetation indices**

- 106 2.3.1. Water Supply Vegetation Index (WSVI):
- $107 \quad WSVI = NDVI/T_s \tag{1}$
- 108 Where

109  $T_s$  is the brightness temperature channel or related remote sensing imagery estimated. The 110 smaller this index is, the more severe the drought is.

111 2.3.2. Soil Adjusted Vegetation Index (SAVI):

112 
$$SAVI = \frac{(NIR-R)}{(NIR+R)*(1+L)}$$
(2)

- 113 Where,
- 114 *NIR* is the Near Infrared band
- 115 *R* is the Red band
- 116 *L* is the soil brightness correction factor, commonly L = 0.5, (Huete, 1988).
- 117
- 118



(3)



## 119 2.3.3. Moisture Stress Index (MSI):

120 
$$MSI = \frac{SWIR_1}{NIR}$$

- 121 Where
- 122  $SWIR_1$  is the Short-wave Infrared band 1
- 123 *NIR* is the Near Infrared band
- 124 2.3.4. Normalized Difference in Level Index (NDII):

125 
$$NDII = \frac{(NIR - SWIR_1)}{(NIR + SWIR_1)}$$
 (4)

- 126 Where
- 127 *NIR* is the Near Infrared band
- 128  $SWIR_1$  is the Short-wave Infrared band 1.

129

# 130 **2.4. Estimation of soil salinity index**

131 Soil salinity indices are principally adjusted to detect salt mineral in soils based on the different

132 responses of salty soils to various spectral bands. The following equation to map soil salinity was

used after Elhag (2016):

$$134 \quad SI = (G \times R)/B \tag{5}$$

- 135 Where,
- 136 *B* is the Blue band
- 137 G is the Green band
- 138 *R* is the Red band





### 139 2.5. Regression Analyses

- 140 The purpose of the regression analyzes is to envisage the regression potentials between soil
- salinity index from one side and the rest of the hydrological drought indices from the other side.
- 142 Principle Component Analysis (PCA) and Artificial Neural Network (ANN) were the
- 143 implemented approaches. The PCA is to transform a set of likely correlated with unlikely
- 144 correlated variables. Principal components number is less/equal to the variables original number.
- 145 Following Lorenz (1956), PCA fundamental equations are:
- 146 First vector W (1) has to be answered as following:

147 
$$w_{(1)} = \arg \max_{\|w\|=1} \left\{ \sum_{i} (t_1)_{(i)}^2 \right\} = \arg \max_{\|w\|=1} \left\{ \sum_{i} (x_i \cdot w)^2 \right\}$$
 (6)

148 The matrix form of the above equation gives the following:

149 
$$w_{(1)} = \arg \max_{\|w\|=1} \{ \|Xw\|^2 \} = \arg \max_{\|w\|=1} \{ w^T X^T Xw \}$$
 (7)

150  $W_{(1)}$  has to be answered as following:

151 
$$w_{(1)} = \arg \max\left\{\frac{w^T x^T x w}{w^T w}\right\}$$
 (8)

Originated  $w_{(1)}$  suggests that first component of a data vector  $x_{(i)}$  can then be expressed as a score of  $t_{1(i)} = x_{(i)} \cdot w_{(1)}$  in the transformed co-ordinates, or as the corresponding vector in the original variables,  $\{x_{(i)} \cdot w_{(1)}\} w_{(1)}$ .

155 The neural network regression model is written as:

156 
$$Y = \alpha + \sum_{h} w_{h} \phi_{h} (\alpha_{h} + \sum_{i=1}^{p} w_{ih} X_{i})$$
 (9)

157 Where

158 Y = E(Y|X). This neural network model has 1 hidden layer but it is possible to have additional 159 hidden layers.





160 The  $\phi(z)$  function used is hyperbolic tangent activation function. It is not activation

161 for the hidden layers.

162 
$$\phi(z) = \tanh(z) = \frac{1 - e^{-2z}}{1 + e^{-2z}}$$
 (10)

163 It is significant that the final outputs to be linear not to constrain the predictions to be between 0 164 and 1. A simple diagram of a skip-layer neural network is illustrated in Figure 2. The equation 165 for the skip-layer neural network for regression is shown below.

166 
$$Y = \alpha + \sum_{i=1}^{p} \beta_i X_i + \sum_h w_h \phi_h (\alpha_h + \sum_{i=1}^{p} w_{ih} X_i)$$
(11)

167 It should be clear that these models are highly parameterized and thus, will tend to overfit the

training data. Cross-validation is, therefore, critical to make sure that the predictive performance

169 of the neural network model is adequate.

## 170 Figure 2. Artificial Neural Network scheme with 1 hidden layer and 3 nodes.

171 Recall the skip-layer neural network regression model looks like this:

172 
$$Y = \alpha + \sum_{i=1}^{p} \beta_i X_i + \sum_h w_h \phi_h (\alpha_h + \sum_{i=1}^{p} w_{ih} X_i)$$
(12)

However, this model most likely overfits the training data. Consequently, determination of the
adequate performance of the ANNs model is a must. Five different criteria are used: the Pearson
coefficient of correlation (R), the root mean square error (RMSE), the mean absolute Deviation
(MAD), the negative log-likelihood and the unconditional sum of squares (SSE). Basically,
RMSE is the examined parameter for comparability reasons. RMSE can be computed as:

178 
$$RMSE = \sqrt{\frac{1}{T_0}} \sum_{t=1}^{T_0} (y_1 - \dot{y}_1)^2$$
 (13)

Where *t* is the time index,  $\hat{y}_t$  and  $y_t$  are the simulated and measured values. Principally, the higher value of R and smaller values of RMSE ensure the better performance of the model.





#### 181 **3. Results and Discussion**

Realization of the hydrological drought indices was exercised after a comprehensive remote 182 sensing data correction. Basically, atmospheric correction and spatial enhancement were 183 practiced utilizing Landsat 8 data acquired over the designated study area. The four hydrological 184 drought indices were shown in Figures 3 to 6. Stochastic algorithms of WSVI and SAVI 185 186 mapping (Figures 3 and 4) showed spatial coherence with a higher drought indices value within the agricultural area rather than the surrounding (Ceccato et al., 2001; Daughtry et al., 2004). 187 On the contrary, MSI exercised as a deterministic drought index, it's nearly unaffected by 188 changing water content. Conducted results showed two classes of stresses, stressed and no stress. 189 The no stress class wated within the agricultural area and the stressed area we esented along the 190 agricultural peripheral areas (Figure 5) where higher values indicate greater water stress and less 191 water content. This could be explained rationally by the presence of irrigational sprinkles (Hunt 192 193 et al., 1989; Ceccato et al., 2001). NDII is also a stochastic algorithm and was exercised in the current research due to the higher sensitivity of Infrared band to detect change 194 in water content of plant canopies (Hardisky et al., 1983). Spatial distribution of NDII (Figure 6) was mapped 195 196 accordingly with WSVI and SAVI indices, in which higher NDII values methods higher water 197 content (Jackson et al., 2004). There are several algorithms to map soil salinity based on 198 utilization of different remote sensing data as well as different ecological systems. An adequate 199 NDSI algorithm was carried out according to Elhag (2016) findings in arid ecosystems. In Figure 7, NDSI was mapped in the designated study area showed spatial variation of salted soils, 200 especially the new agricultural expansion at the southern west part of the designated study area 201 202 due to the sprinkle movement drove the accumulation of excess waters at the peripherals of the agricultural areas (Lunetta et al., 2002; Konukcu et al., 2006). 203





204	Figure 3. Water Supply Vegetation Index (WSVI) thematic map over the study area.
205	
206	Figure 4. Soil Adjusted Vegetation Index (SAVI) thematic map over the study area.
207	
208	Figure 5. Moisture Stress Index (MSI) thematic map over the study area.
209	
210	Figure 6. Normalized Difference Infrared Index (NDII) thematic map over the study area.
211	
212	Figure 7. Normalized Difference Salinity Index (NDSI) thematic map over the study area.
213	
214	Further statistical analyzes were carried out to construe the correspondences between salted soils
215	and different horological drought indices. Regression analysis demonstrated in Figure 8 showed
216	that salinity increases with lower WSVI and SAVI (Figure 8 a, b) which lained due we salt
217	accumulation in soils. Under salinity stress conditions, there is no enough available water in soils
218	for proper vegetation growth (Lunetta et al., 2002; Yang et al., 2011).
219	Generally, MSI values (Figure 8 c) are high in the study area because of the excess irrigation
220	regime adopted to overcome the high evaporation rates (Elhag and Bahrawi, 2014; Elhag, 2016).
221	Excess irrigation regimes in poor drain soils lead to waterlogging problems and salts accusation
222	(Elhag, 2016).
223	Due to NDII higher sensitivity to water, NDII values increases with higher NDSI values. Salts
224	accumulation caused by excessive irrigation is the driving force behind the proportional
225	increment of NDII values in conjunction with NDSI values phonstrated in Figure 8d (Jackson
226	et al., 2004; Shishi et al., 2015).





227	Figure 8. Regression analyzes pf NDSI (ppm) against horological drought indices.
228	
229	Figure 9 demonstrated the Principal Component Analysis along with the Factor Analysis.
230	Moreover, eigenvalue decomposition is also demonstrated. WSVI and SAVI were grouped
231	together. On the other hand, NDII and MSI were individually plotted against the former indices.
232	
233	Figure 9. Principle Component Analysis.
234	
235	Similar results conducted from the Scatter Plot Matrix and the companion correlation matrix
236	shown in Figure for and Table 1. A high correlation is distinguished between WSVI and SAVI
237	while wative correlation with the between WSVI and SAVI from one side and MSI Ale NDII
238	from the other side.
239	Figure 10. Scatterplot Correlation Matrix.
240	
241	Table 1. Correlation matrix.
242	
243	In Table 2, NDSI regression analysis shows that NDII is the proper fit based on different
244	regression parameters (Rodgers and Nicewander, 1988). Spearman's correlation demonstrated in
245	Table 3 supports PCA results. Hydrological drought indices were classified into two categories,
246	MSI, and NDII in one category and WSVI and SAVI in the other one. The elements of each
247	category are positively correlated. MSI and NDII were significantly correlated; WSVI and SAVI
248	were highly correlated. Moreover, any other combinations of the four hydrological drought
249	indices were not correlated.





The ANN analysis was carried out under 1 hidden layer, 3 nodes, and hyperbolic tangent activation function conditions. These conditions were carefully exercised to prevent the algorithm overfitting, analysis is demonstrated in Table 4. NDII expressed the highest RMSE which indicates that NDSI and NDII are statistically the best fit ((Jiang, 2013). SAVI comes at the second best fit followed by WSVI. MSI failed to fit NDSI values comprehensively like the former hydrological drought indices (Jones and Marshall, 1992; Jiapaer et al., 2011).

- 256 Table 2. Regression analysis.
- 257
- 258 Table 3. Spearman's correlation.
- 259
- 260 Table 4. Neural Network Analysis .
- 261
- 262 **4. Conclusion**

263 The findings of the current research emphasized on the importance of the horological drought indices to envisage the adverse effects of salts accumulation in poorly drained soils. Remote 264 Sensing techniques were satisfactory implement and interpreted in term of soil salinity mapping 265 in consort with hydrological drought indices. Normalized Difference Infrared Index was 266 267 statistically proved to be the Normalized Difference Salinity Index profound, followed by Soil Adjusted Vegetation Index and Water Shortage Vegetation Index respectively. Principal 268 Component Analysis and Artificial Neural Network Analysis are complementary tools to 269 understand the regression pattern of the hydrological drought indices in the designated study 270 area. Further work needs to be considered towards the restrictiveness of the drastic effect of salts 271 accumulation within the study area. 272





#### 273 Acknowledgment

- This work was supported by the Deanship of Scientific Research (DSR), King Abdulaziz University,
  Jeddah, under grant No. (155-36-1437D). The authors, therefore, gratefully acknowledge the DSR
  technical and financial support.
- 277

278	References   O
279	Al-Zahrani, K. H. and Baig, M. B. (2011). Water in the Kingdom of Saudi Arabia: Sustainable
<mark>280</mark>	Management Options. The Journal of Animal and Plant Sciences, 21: 601–604.

- Bannari, A., Guedona, A. M., El-Hartib, A., Cherkaouic, F. Z. and El-Ghmari, A. (2008).
  Characterization of Slightly and Moderately Saline and Sodic Soils in Irrigated
  Agricultural Land using Simulated Data of Advanced Land Imaging (EO-1) Sensor.
  Communications in Soil Science and Plant Analysis, 39(19-20): 2795-2811.
- Boegh, E., Soegaard, H., Broge, N., Hasager, C. B., Jensen, N. O., Schelde, K., and Thomsen, A.
  (2002). Airborne multispectral data for quantifying leaf area index, nitrogen
  concentration, and photosynthetic efficiency in agriculture. Remote Sensing of
  Environment, 81: 179–193.
- Bouman B. A. M. and Tuong, T. P. (2001). Field water management to save water and increase
  its productivity in irrigated rice. Agricultural Water Management, 49(1):11-30.
- Box, G. E. P. (1954). Some Theorems on Quadratic Forms Applied in the Study of Analysis of
  Variance Problems, II: Effects of Inequality of Variance and Correlation Between Errors
  in the Two-Way Classification. Annals of Mathematical Statistics, 1: 69-82.
- Broge, N. H., and Mortensen, J. V. (2002). Deriving green crop area index and canopy
  chlorophyll density of winter wheat from spectral reflectance data. Remote Sensing of
  Environment, 81: 45–57.
- 297 Ceccato, P., Flasse, S., Tarantola, S., Jacquemoud, S. and Gregoire, J. M. (2001). Detecting
  298 vegetation leaf water content using reflectance in the optical domain. Remote Sensing of
  299 Environment, 77: 22–33.
- Clevers, J. G. P. W. (1989). The application of a weighted infrared-red vegetation index for
  estimating leaf-area index by correcting for soil-moisture. Remote Sensing of
  Environment, 29: 25–37.





303 Colombo, R., Bellingeri, D., Fasolini, D. and Marino, C. M. (2003). Retrieval of leaf area index 304 in different vegetation types using high resolution satellite data. Remote Sensing of Environment, 86: 120-131. 305 Curran, P. J. (1983a). Estimating green LAI from multispectral aerial-photography. 306 Photogrammetric Engineering and Remote Sensing, 49: 1709–1720. 307 Curran, P. J. (1983b). Multispectral remote-sensing for the estimation of green leaf area index. 308 Philosophical Transactions of the Royal Society of London Series a Mathematical 309 310 Physical and Engineering Sciences, 309: 257–270. Daughtry, C. S. T., Hunt Jr., E. R. and McMurtrey III, J. E. (2004). Assessing crop residue cover 311 using shortwave infrared reflectance. Remote Sensing of Environment 90, 126-134. 312 Douaoui, A. K., Herve', N. and Walter, C. (2006). Detecting salinity hazards within a semiarid 313 context by means of combining soil and remote sensing data. Geodema, 134: 217-230. 314 Elhag, M. (2016). Evaluation of Different Soil Salinity Mapping Using Remote Sensing 315 Techniques in Arid Ecosystems, Saudi Arabia. Journal of Sensors, 2016: 96175-96175. 316 Elhag, M. (2014a). Remotely Sensed Vegetation Indices and Spatial Decision Support System 317 for Better Water consumption Regime in the Nile Delta. A Case Study for Rice 318 319 Cultivation Suitability Map. Life Science Journal, 11(1): 201-209. Elhag, M. (2014b). Sensitivity Analysis Assessment of Remotely Based Vegetation Indices to 320 321 Improve Water Resources Management. Environment Development and Sustainability, 16(6): 1209-1222. 322 Elhag, M. and Bahrawi, J. (2014). Conservational Use of Remote Sensing Techniques for a 323 324 Novel Rainwater Harvesting in Arid Environment. Environmental Earth Sciences, 325 72(12): 4995-5005. Gobron, N., Pinty, B., Verstraete, M. M. and Widlowski, J. L. (2000). Advanced vegetation 326 indices optimized for up-coming sensors: Design, performance, and applications. IEEE 327 328 Transactions on Geoscience and Remote Sensing, 38: 2489-2505. Govaerts, Y. M., Verstraete, M. M., Pinty, B. and Gobron, N. (1999). Designing optimal spectral 329 indices: A feasibility and proof of concept study. International Journal of Remote 330 Sensing, 20: 1853-1873. 331





- Hardisky, M., Klemas, V. and Smart, R. (1983). The influence of soil salinity, growth form, and
  leaf moisture on the spectral radiance of *Spartina alterniflora* canopies. Photogrammetric
  Engineering & Remote Sensing, 49: 77–83.
- Huete, A., Didan, K., van Leeuwen, W., Miura, T. and Glenn, E. (2008). MODIS vegetation
  indices. In Land Remote Sensing and Global Environmental Change: NASA's Earth
  Observing System and the Science of ASTER and MODIS, 125 146.
- Hunt, Jr. E.R., and Rock, B.N. (1989). Detection of changes in leaf water content using near- and
  middle infrared reflectances. Remote Sensing of Environment, 30: 43–54.
- Jackson, T. J., Chen, D. Y., Cosh, M., Li, F. Q., Anderson, M., Walthall, C., Doriaswamya, P.
  and Hunt, Jr. R. E. (2004). Vegetation water content mapping using Landsat data derived
  normalized difference water index for corn and soybeans. Remote Sensing of
  Environment, 92, 475–482.
- Jiang, B. (2013). Head/Tail Breaks: A New Classification Scheme for Data with a Heavy-Tailed
   Distribution. The Professional Geographer, 65(3): 482-494.
- Jiapaer, G., Chen, X. and Bao, A. M. (2011). A comparison of methods for estimating fractional
  vegetation cover in arid regions. Agricultural and Forest Meteorology, 151(12): 1698–
  1710.
- Jiapaer, G., Chen, X. and Bao, A. M. (2011). A comparison of methods for estimating fractional
  vegetation cover in arid regions. Agricultural and Forest Meteorology, 151(12): 1698–
  1710.
- Jones, R. and Marshall, G. (1992). Land salinization, waterlogging and the agricultural benefits
   of a surface drainage scheme in Benerembah irrigation district. Review of Marketing and
   Agricultural Economics, 60: 173–189.
- Kaufman, Y.J. and Tanré, D. (1992). Atmospherically resistant vegetation index (ARVI) for
   EOS-MODIS. Transactions on Geoscience and Remote Sensing, 30: 261–270.
- Kerr, J. and Ostrovsky, M. (2003). From space to species: ecological applications for remote
   sensing. Trends in Ecology and Evolution, 18: 299-305.
- Konukcu, F., Gowing, J. W. and Rose, D. A. (2006). Dry drainage: a sustainable solution to
   (waterlogging and salinity problems in irrigation areas? Agricultural Water Management
- **361 83**: 1–12,





362 Konukcu, F., Gowing, J.W. and Rose, D.A. (2006). Dry drainage: A sustainable solution to 363 waterlogging and salinity problems in irrigation areas? Agricultural Water Management, 83: 1-12. 364 Leprieur, C., Kerr, Y. H., Mastorchio, S. and Meunier, J. C. (2000). Monitoring vegetation cover 365 across semi-arid regions: Comparison of remote observations from various scales. 366 International Journal of Remote Sensing, 21(2):281-300. 367 Lorenz, E. N. (1956). Empirical orthogonal functions and statistical weather prediction. MIT 368 Department of Meteorology, Statistical Forecast Project Rep. 1, 49 pp. [Available from 369 370 Dept. of Meteorology, MIT, Massachusetts Ave., Cambridge, MA 02139. 371 Montandon, L. M. and Small, E. E. (2008). The impact of soil reflectance on the quantification of the green vegetation fraction from NDVI. Remote Sensing Environment, 112: 1835 -372 373 1845. 374 Moulin, S. and Guerif, M. (1999). Impacts of model parameter uncertainties on crop reflectance 375 estimates: A regional case study on wheat. International Journal of Remote Sensing, 20: 213-218. 376 Pettorelli, N., Vik, J., Mysterud, A., Gaillard, J., Tucker, C. and Stenseth, N. (2005). Using the 377 378 satellite-derived NDVI to assess ecological responses to environmental change. Trends in 379 Ecology and Evolution, 20: 503-510. Pinty, B. abd Verstraete, M. M. (1992). GEMI: A non-linear index to monitor global vegetation 380 from satellites. Plant Ecology, 101: 15-20. 381 382 Pinty, B., Lavergne, T., Widlowski, J. L., Gobron, N. and Verstraete, M. M. (2009). On the need to observe vegetation canopies in the near-infrared to estimate visible light absorption. 383 Remote Sensing of Environment, 113: 10-23. 384 Psilovikos, A. and Elhag, M. (2013). Forecasting of Remotely Sensed Daily Evapotranspiration 385 386 Data over Nile Delta Region, Egypt, Water Resources Management, 27(12): 4115-4130. Qi, J., Chehbouni, A., Huete, A. R.; Kerr, Y. H. and Sorooshian, S. (1994). A modified soil 387 388 adjusted vegetation index. Remote Sensing of Environment, 48: 119-126. Rodgers, J. L. and Nicewander, W.A. (1988). Thirteen ways to look at the correlation coefficient. 389 The American Statistician, 42 (1): 59-66. 390





391	Verstraete, M. M., Pinty, B. and Myneni, R. B. (1996). Potential and limitations of information
392	extraction on the terrestrial biosphere from satellite remote sensing. Remote Sensing of
393	Environment, 58: 201–214.
394	Wardlow, B. D. and Egbert, S. L. (2010). A comparison of MODIS 250-m EVI and NDVI data
395	for crop mapping in the U.S. Central Great Plains. International Journal of Remote
396	Sensing, 31(3): 805-830.
397	Xiao, X., He, L., Salas, W., Li, C., Moore, B., Zhao, R., Frolking, S. and Boles, S. (2002).
398	Quantitative relationships between field-measured leaf area index and vegetation index
399	derived from vegetation images for paddy rice fields. International Journal of Remote
400	Sensing, 23: 3595–3604.
401	Yang, J. Y., Zheng, W., Tian, Y., Wu, Y. and Zhou, D. W. (2011). Effects of various mixed salt-
402	alkaline stresses on growth, photosynthesis, and photosynthetic pigment concentrations
403	of Medicago ruthenica seedlings. Photosynthetica, 49: 275-284.
404	
405	
406	
407	
408	
409	
410	
411	
412	
413	
414	
415	
416	
417	
418	
419	
420	
421	













432

Figure 2. Artificial Neural Network scheme with 1 hidden layer and 3 nodes. 433

434







436

437 Figure 3. Water Supply Vegetation Index (WSVI) thematic map over the study area.

438







440

441 Figure 4. Soil Adjusted Vegetation Index (SAVI) thematic map over the study area.

442









446

444







448

449 Figure 6. Normalized Difference Infrared Index (NDII) thematic map over the study area.

450







453 Figure 7. Normalized Difference Salinity Index (NDSI) thematic map over the study area.



























486

# 487 **Table 1. Correlation matrix.**

		NDII	MSI	SAVI	WSVI
	NDII	1	0.7182080406	-0.708975719	-0.703572559
	MSI		1	-0.888156103	-0.88249756
	SAVI			1	0.9977255509
	WSVI				1
488					
489					
490					
491					
492					
493					
494					
495					
496					
497					
498					
499					
500					
501					
502					
503					
504					
505					
506					





# 507 Table 2. Regression analysis.

		NDII	MSI	SAVI	WSVI
	RSquare	0.798566127	0.254999657	0.246131379	0.243463225
	RSquare Adj	0.797205088	0.249965871	0.241037672	0.23835149
	<b>Root Mean Square Error</b>	31.88199207	0.384262574	0.202130562	0.000447112
	Mean of Response	124.5466667	0.733333333	0.286361262	0.000611978
	<b>Observations (Sum Wgts)</b>	150	150	150	150
508					
500					
509					
510					
511					
512					
513					
514					
515					
516					
510					
517					
540					
518					
519					
520					
521					
521					
522					
F 2 2					
525					
524					
525					
526					
527					
528					
J20					





# 529 Table 3. Spearman's correlation.

	Variable	By Variable	Correlation	Count	Lower 95%	Upper 95%	Significance probability
	MSI	NDII	0.7182	150	0.6305	0.7878	*
	SAVI	NDII	-0.7090	150	-0.7805	-0.6191	NS
	SAVI	MSI	-0.8882	150	-0.9178	-0.8487	NS
	WSVI	NDII	-0.7036	150	-0.7763	-0.6124	NS
	WSVI	MSI	-0.8825	150	-0.9136	-0.8412	NS
	WSVI	SAVI	0.9977	150	0.9969	0.9984	**
53	0      * is sig	nificant, ** is h	ighly significa	nt, NS is	non-significan	t	
БЭ	1						
55	T						
53	2						
	•						
53	3						
53	4						
	_						
53	5						
53	6						
	_						
53	7						
53	8						
53	9						
54	0						
	-						
54	1						
54	2						
5-4	-						
54	3						





# 544 Table 4. Neural Network Analysis.

		T	X7-12-1-42 N/		
	DC	1 raining Measures	validation Measures	_	
	RSquare	0.7574526	0.6698156		0.6
_	RMSE	0.0999530	0.0972931	- 132,882	0.4
II	Mean Abs Dev	0.0571881	0.0436599	ā	
Z	<ul> <li>LogLikelihood</li> </ul>	-88.411680	-45.554430	2	0.2
	SSE	0.9990600	0.4732975		
	Sum Freq	100	50		
	RSquare	0.3032101	0.0893892	0.82851/	0.6-1 \
	RMSE	0.2388872	0.1869959	ž	0.4
10	Mean Abs Dev	0.1203075	0.0628425		0.2
Ž	-LogLikelihood	-1.2825260	-12.886510		°]
	SSE	5.7067096	1.7483727		0.8
	Sum Freq	100	50	_ 0.250643	0.6
	~			SA	0.4
	RSquare	0.7565419	0.6698155		0.2
	RMSE	0.1499295	0.1459397		· · · · · · · · · · · · · · · · · · ·
M	Mean Abs Dev	0.0857822	0.0654899		0.002
SA	-LogLikelihood	-47.865170	-25.28115	= 0.000538	0.0015
•1	SSE	2.2478847	1.0649203	S.	0.001
	Sum Freq	100	50	_	0.0005-
	1				0-1
	RSquare	0.7533827	0.6619429		
	RMSE	0.0003280	0.0003226		0 2 0 2 0 0
<b>I</b>	Mean Abs Dev	0.0001876	0.0001451		00004
MS N	-LogLikelihood	-660.35100	-331.01460		28104
	SSE	1.08E-05	5.20E-06		Soil
	Sum Freq	100	50		Salinity

545

546