

Interactive comment on “A comparison of gap-filling algorithms for eddy covariance fluxes and their drivers” by Atbin Mahabbati et al.

Atbin Mahabbati et al.

atbin.m@hotmail.com

Received and published: 10 December 2020

GENERAL COMMENTS

RC02: The paper presents a detailed evaluation of eight algorithms for gap-filling time series data, using eddy covariance data as a target for the comparisons. The content about the algorithms and the metrics for comparisons are a strong feature of the paper. However, it is more limited in advancing the knowledge of best practices for eddy covariance and micro-meteorological data gap-filling. In other words, the evaluation of the algorithms against each other is of interest, but the chosen test domain is not clearly impacted. It seems that to really benefit the knowledge of methods for gap-filling eddy covariance data, longer time series and more representative gap scenarios would

[Printer-friendly version](#)

[Discussion paper](#)



be necessary, as well as a clear comparison to more established methods. Multi-year datasets are key to properly evaluate these algorithms. Such datasets are now widely available, so it is unclear why only 2013 was used. With this aspect in mind, it seems clear that in the evaluation of the first objective of the paper longer gaps led to disproportional increases in uncertainty. This might not have happened if other years without gaps for the same season were available, for instance. More direct comparisons to "classic" gap-filling algorithms would have helped in this evaluation. Implementations of algorithms such as MDS are now widely available, including as part of OzFlux's own OzFluxQC software package. The comparison of newer methods is informative, but unless compared to currently used solutions, it's hard to assess the improvement. Although the authors are correct, and performance of the MDS algorithm was shown to be comparable to ANNs before, parameterizing MDS is much simpler (no choices in layers, nodes, iterations, or window sizes) and would lead to a more robust and clear comparison.

AC: Please note that as a PhD student whose thesis is based on a series of papers, the current paper is the very first one that has mainly provided as the initial attempt to find out how different algorithms would perform against each other. As such, almost all the points mentioned in the general comment, which are helpful, would be covered in the second paper, e.g. including multiple-year datasets, and applying different random gap scenarios. However, as the second referee has mentioned, we accept the idea of adding the results of the MDS in the current study. Last but not least, the year 2013 was chosen for the fact that the data during the period had less missing data, and that year was a common year of available data amongst all five towers that their data were used. Besides, most of the researches have been done in the field includes just one or two years of data, so the results of this paper can be compared with the majority of similar previous researches. Besides, many researchers still fill the annual gaps by using only the data of that year, thus using a year of data for training the algorithms can still be justified.

[Printer-friendly version](#)[Discussion paper](#)

RC02: Should the authors choose to really focus on the comparison among the methods presented, I would suggest adding all the comparison metrics RMSE, R2, MBE, etc., for all sites individually and combinations thereof as supplementary materials, making this a valuable and thorough comparison of methods, and reducing the focus from the application to eddy covariance. If the intention really is to show the impact on EC, longer time series and more direct comparisons to current methods would be necessary.

AC: We are happy with adding all the comparison metrics for all sites as supplementary materials. Besides, the intention was to make a comparison between different algorithms, and as such, in case using a year of data is insufficient, it would be equally insufficient for all algorithms.

SPECIFIC COMMENTS

RC02: On the ancillary datasets, it seems they introduce some entanglement to this evaluation. One of the key advantages of purely empirical methods, such as the ones presented in the paper, is that they will not be biased by predefined models (like the reanalysis datasets) or atmospheric interferences (like the MODIS data). After an evaluation without datasets such as these, adding them to improve the methods would be a natural choice. However, without the unbiased evaluation it is hard to qualify the sources of uncertainty in the paper's evaluation.

AC: Even though this is true, the ancillary data used in the current study have been used to gap-fill the drivers' data, and not the fluxes directly. As such, it might not be a concern. Nonetheless, since the corresponding author is going to use the EVI data as one of the gap-filling features for Fc, the corresponding author wonders whether the referee has any suggestion to address the issue?

RC02: Although the performance criteria selected for the paper work well, it is curious to see that the methods all seem to represent high variabilities but fail to capture the extremes, as the authors point out for CO2 and latent heat fluxes – and this doesn't

[Printer-friendly version](#)[Discussion paper](#)

seem to be the case for sensible heat flux. Could this be an issue of the underlying data requiring further quality control before the gap-filling methods are applied? Or maybe this is an artifact of the period selected in the examples?

AC: This was one of the surprising things raised during the study, and to be honest, we do not have a solid answer to that yet. However, estimating the sensible heat flux is an easier task as against the two others. This can justify the exception of sensible heat flux. For F_c , and F_e , our best guess is that the issue happens due to lack of information (hidden features). We will try to figure that out in the second paper of this series.

RC02: The following claim requires either more details or a reference, otherwise it's not possible to know what concerns/challenges the authors are referring to and what aspects of gap-filling the paper is aiming to address: "...there are some serious concerns regarding the challenges associated with the technique, e.g. data gaps and uncertainties."

AC: That is right. The relevant references, as well as more explanation, will be added to clarify the sentence.

RC02: The $\pm 25 \text{gCm}^{-2}\text{y}^{-2}$ (Moffat et al. 2007) and $\pm 30 \text{gCm}^{-2}\text{y}^{-2}$ (Richardson & Hollinger 2007) are dependent on the underlying datasets used for the evaluation. These numbers should not be taken as general benchmarks.

AC: That is the right point. This part will be modified, and an appropriate reference will be added.

RC02: In the sentence "Nevertheless, one of the concerns regarding this algorithm is that the independent variables, here meteorological drivers, might be auto-correlated." it is unclear why this would be a concern, since the meteorological drivers being auto-correlated is one of the assumptions that allow the MDS method to work.

AC: The comment is true. We will delete the sentence.

RC02: The sentence "This challenge becomes acute when the gaps happen within

a period when the ecosystem behaviour is changing and thereby showing different response under similar meteorological conditions." is another reason why multi-year datasets should be used to compare these algorithms.

AC: As mentioned earlier, we used one year of data because: (a) the focus had been on algorithm comparison, (b) most previous researches used a year or two, so our results could be more comparable with them, and (c) 2013 was the year during which the datasets for all five towers were smaller proportions of gaps. Finally, in the second paper of the series, we are using up to five years of data for training and testing. Hence, the concern would be considered in the bigger picture.

RC02: The gap scenarios and training windows selected are highly structured and rigid. It's unclear how the evaluation over these scenarios would translate into real-world examples, which have both structured gaps (e.g., from sensor failures) and arbitrary gaps (e.g., from data filtering). It seems it would be important to use at least one scenario with gaps and training data both randomized, and also combinations of lengths for gap windows and training windows.

AC: This is a good and constructive suggestion. However, please note that this paper is the first one of a series of papers the corresponding author has to write down as his PhD thesis. For the second one, which the corresponding author is working on right now, randomly selected gaps have been superimposed.

TECHNICAL CORRECTIONS

Abstract -

RC02: The acronyms RF and CLR were referenced before being defined

AC: Thank you for letting us know that. Those acronyms would be predefined in the revised version.

RC02: - "...RF provided more consistent results with less bias, relatively." It would be clearer to describe "relatively" to what in this sentence.

AC: That is a helpful suggestion. The authors mean related to the rest of the algorithms used in the study. The sentence would become modified accordingly.

RC02: - This sentence is a bit unclear "In each scenario, the gaps covered the data for the entirety of 2013 by consecutively repeating them, where, in each step, values were modelled by using earlier window data." Were measured and modelled data used simultaneously in training? â€” Introduction

AC: The reviewer has pointed out an important issue that is the explanation of the scenarios is not clear enough, particularly because this was the issue for the first reviewer as well. As such, we need to rewrite the sentence to make sure that the method is clear enough and easy to understand.

RC02: - "...and not measured at the point." Maybe could be "not measured at a point scale"?

AC: That is right. We will correct it.

RC02: - A more classic reference for FLUXNET is: Baldocchi et al. 2001. FLUXNET: A New Tool to Study the Temporal and Spatial Variability of Ecosystem-Scale Carbon Dioxide, Water Vapor, and Energy Flux Densities. BAMS, 11: 2415-2434.

AC: Thank you for reminding that reference. We would use include it in the introduction.

RC02: - And more appropriate references for EUROFLUX and AmeriFlux are: Aubinet, M. et al. 1999. Estimates of the Annual Net Carbon and Water Exchange of Forests: The EUROFLUX Methodology. Advances in Ecological Research, pp. 113–175. Law, B. 2007. AmeriFlux Network aids global synthesis. Eos, 88, 286–286. Novick, K. A. et al 2018. The AmeriFlux network: A coalition of the willing. AFM, 249:444-456.

AC: We would consider the references.

RC02: - "Despite the capability of EC to frequently validate process modelling analyses..." might be more precisely phrased as something like "Despite EC data being

[Printer-friendly version](#)[Discussion paper](#)

frequently used to validate process modelling analyses..."

AC: Yes. That would make the sentence more natural and smooth.

RC02: - "[...] Moffat et al. (2007) compared a couple of different commonly-used gap-filling algorithms"; in fact, Moffat et al. 2007 compared 15 gap-filling techniques.

AC: Yes, we would mention the exact number, instead of "a couple".

- Materials and Methods

RC02: - "and Tumberumba form 2011 to 2013..." form -> from

AC: Thank you for mentioning the mistake.

RC02: - "Each algorithm was tuned up individually using gird search,..." gird -> grid

AC: Thank you for mentioning the mistake.

âĀĤ Results

RC02: - Even with a maximum zoom in the PDF file, it is rather hard to read the axis for Figures 3 and 4

AC: That happens because the size/resolution of the figures is not big/high enough. The figures would be replaced with appropriate ones.

âĀĤ Discussion

RC02: - This sentence is unclear: "That is because ANNs have been checking out for a long time in different locations and considered as one of the most reliable algorithms in the field for more than a decade"

AC: The authors mean occasional superiority of random forest algorithm, needs to be happen in several future researches to convince us to suggest RF instead of ANNs, or identify the algorithm as another standard method. We will add a sentence to clarify the point.

[Printer-friendly version](#)

[Discussion paper](#)



Interactive comment on Geosci. Instrum. Method. Data Syst. Discuss.,
<https://doi.org/10.5194/gi-2020-21>, 2020.

GID

Interactive
comment

[Printer-friendly version](#)

[Discussion paper](#)

